

UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

Docket No.
51271/35:1

Total Pages in this Submission

(Only for new nonprovisional applications under 37 CFR 1.53(b))

TO THE COMMISSIONER FOR PATENTS

Mail Stop Patent Application
P.O. Box 1450
Alexandria, VA 22313-1450

COPY

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

ADENO-ASSOCIATED VECTOR COMPOSITIONS FOR EXPRESSION OF FACTOR VIII

and invented by:

Linda B. Couto
Peter C. Colosi
Xiaobing Qian

If a CONTINUATION APPLICATION, check appropriate box and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: 09/740,211

Which is a:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: 09/470,618

Which is a:



☐ Continuation ☐ Divisional ☒ Continuation-in-part (CIP) of prior application No.: 09/364,862

☒ Provisional Application Serial Nos. 60/125,974 and 60/104,994

Enclosed are:

Application Elements

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification having sixty-three (63) pages and including the following:
 - a. ☒ Descriptive Title of the Invention
 - b. ☒ Cross References to Related Applications (if applicable)
 - c. ☐ Statement Regarding Federally-sponsored Research/Development (if applicable)
 - d. ☒ Reference to Sequence Listing, a Table, or a Computer Program Listing Appendix
 - e. ☒ Background of the Invention
 - f. ☒ Brief Summary of the Invention
 - g. ☒ Brief Description of the Drawings (if filed)
 - h. ☒ Detailed Description
 - i. ☒ Claim(s) as Classified Below
 - j. ☒ Abstract of the Disclosure

CERTIFICATE OF MAILING BY "EXPRESS MAIL" (37 CFR 1.10) Applicant(s): Couto et al.			Docket No. 51271/35:1	
Serial No. Not Yet Assigned	Filing Date August 1, 2003	Examiner Not Yet Assigned		Group Art Unit Not Yet Assigned
Invention: ADENO-ASSOCIATED VECTOR COMPOSITIONS FOR EXPRESSION OF FACTOR VIII				
<div style="float: left; width: 150px;">  </div> <div style="float: right; font-size: 4em; transform: rotate(-15deg); opacity: 0.5;">COPY</div> <div style="clear: both;"></div>				
I hereby certify that the following correspondence:				
Continuation Patent Application (including the items listed below)				
<i>(Identify type of correspondence)</i>				
is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on				
<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center; margin-right: 20px;"> <u>Aug 1, 2003</u> <i>(Date)</i> </div> </div>				
<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center; margin-right: 20px;"> Mara K. Featherstone <i>(Typed or Printed Name of Person Mailing Correspondence)</i> </div> </div>				
<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center; margin-right: 20px;">  <i>(Signature of Person Mailing Correspondence)</i> </div> </div>				
<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center; margin-right: 20px;"> EU719410029US <i>("Express Mail" Mailing Label Number)</i> </div> </div>				

Note: Each paper must have its own certificate of mailing.

Transmitted:

- Continuation Patent Application (63 pgs.)
- Formal Drawing Sheets (14)
- Verification Under 37 C.F.R. 1.821(e)-(g)
- Utility Patent Application Transmittal Letter (4 pgs.)
- PTO-Credit Card Form 2038 charging the amount of \$415.00
- Assignment including Recordation Form Cover Sheet (2 pgs.)
- Declaration for Patent Application (1 pg.)
- Power of Attorney (1 pg.)
- Verified Statement Claiming Small Entity Status (1 pg.)
- Certificate of Mailing by Express Mail Label No. EU719410029US
- Postcard

THE UNITED STATES PATENT AND TRADEMARK OFFICE:
PLEASE **OPEN** AND RETURN. THANK YOU.

Submitted: Continuation Patent Application (63 pgs.); Formal Drawing Sheets (14); Verification Under 37 C.F.R. 1.821(e)-(g); Utility Patent Application Transmittal Letter (4 pgs.); PTO-Credit Card Form 2038 charging the amount of \$415.00; Assignment including Recordation Form Cover Sheet (2 pgs.); Declaration for Patent Application (1 pg.); Power of Attorney (1 pg.); Verified Statement Claiming Small Entity Status (1 pg.); Certificate of Mailing by Express Mail Label No. EU719410029US; Postcard

Applicant: Couto et al.
Title: ADENO-ASSOCIATED VECTOR COMPOSITIONS
FOR EXPRESSION OF FACTOR VIII

Mailed: August 1, 2003
Docket No.: 51271/35.1 US

08306 U.S. PTO
10/632645
08/01/03

COPY

87

**EXPRESS
MAIL**

POST OFFICE TO ADDRESSEE



* E U 7 1 9 4 1 0 0 2 9 U S *

POSTAL USE ONLY

Day of Delivery <input type="checkbox"/> Next <input type="checkbox"/> Second	Flat Rate Envelope <input type="checkbox"/>
Year: <input type="checkbox"/> 12 Noon <input type="checkbox"/> 3 PM	Postage \$ <u>17.00</u>
Military <input type="checkbox"/> PM <input type="checkbox"/> 2nd Day <input checked="" type="checkbox"/> 3rd Day	Return Receipt Fee
Int'l Alpha Country Code	COD Fee Insurance Fee
Acceptance Clerk Initials	Total Postage & Fees \$ <u>17.00</u>
<input type="checkbox"/> Holiday	

SEE REVERSE SIDE FOR
SERVICE GUARANTEE AND
INSURANCE COVERAGE LIMITS

☐ **WAIVER OF SIGNATURE** (Domestic Only) Conditional merchandise insurance is void if waiver of signature is requested. I wish delivery to be made without obtaining signature of addressee or addressee's agent (if delivery employee judges that article can be left in secure location) and I authorize that delivery employee's signature constitutes valid proof of delivery.

NO DELIVERY ☐ Weekend ☐ Holiday

Customer Signature

Customer Copy
Label 11-B June 2002


USER USE ONLY

PAYMENT Corporate Acct. No. <u>841086</u> <u>51271/35</u>	Federal Agency Acct. No. or Postal Service Acct. No.										
PLEASE PRINT PHONE <u>801</u> <u>578-6992</u>	TO: (PLEASE PRINT) PHONE ()										
NARA K. FEATHERSTONE STOEL RIVES LLP ONE UTAH CENTER 201 SOUTH MAIN STREET, SUITE 1100 SALT LAKE CITY, UT 84111	MAIL STOP PATENT APPLICATION COMMISSIONER FOR PATENTS P.O. BOX 1450 ALEXANDRIA, VA 22313-1450										
ZIP + 4 <table border="1"><tr><td>2</td><td>2</td><td>3</td><td>1</td><td>3</td><td>+</td><td>1</td><td>4</td><td>5</td><td>0</td></tr></table>	2	2	3	1	3	+	1	4	5	0	
2	2	3	1	3	+	1	4	5	0		

FOR PICKUP OR TRACKING CALL 1-800-222-1811

www.usps.com





UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPL NO.	FILING OR 371 (c) DATE	ART UNIT	FIL FEE REC'D	ATTY. DOCKET NO	DRAWINGS	TOT CLMS	IND CLMS
10/632,645	08/01/2003	1632	375	51271/35:1	14	11	1

35735
STOEL RIVES LLP
201 SOUTH MAIN STREET, SUITE 1100
SALT LAKE CITY, UT 84111

CONFIRMATION NO. 5318

FILING RECEIPT



OC0000001602322

Date Mailed: 05/13/2005

Receipt is acknowledged of this regular Patent Application. It will be considered in its order and you will be notified as to the results of the examination. Be sure to provide the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION when inquiring about this application. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please mail to the Commissioner for Patents P.O. Box 1450 Alexandria Va 22313-1450. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections (if appropriate).

Applicant(s)

Linda B. Couto, Pleasanton, CA;
Peter C. Colosi, Alameda, CA;
Xiaobing Qian, Alameda, CA;

Power of Attorney:

Kevin Laurence-38219
John Thompson-40842
Kenneth Chahine-42398
Mary Thomson-43190
Kory Christensen-43548

Scott Young-45249
Mara Featherstone-50390
Matthew Thayne-52280

Domestic Priority data as claimed by applicant

This application is a CON of 09/740,211 12/18/2000 ABN
which is a CON of 09/470,618 12/22/1999 PAT 6,200,560
which is a CIP of 09/364,862 07/30/1999 PAT 6,221,349
which claims benefit of 60/125,974 03/24/1999
and claims benefit of 60/104,994 10/20/1998

Foreign Applications

If Required, Foreign Filing License Granted: 05/13/2005

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is **US10/632,645**

references in the specification to the drawing figures to correspond with any relabeled drawing figures, is required. A copy of the drawing figures showing the proposed changes in red ink should accompany with any drawing changes. Such amendment and/or correction to the drawing figures, if necessary, should be by way of preliminary amendment submitted prior to the first Office action to avoid delays in the prosecution of the application.

Replies should be mailed to: Mail Stop Missing Parts
Commissioner for Patents
P.O. Box 1450
Alexandria VA 22313-1450

A copy of this notice MUST be returned with the reply.

N. Mohammed

Office of Initial Patent Examination (703) 308-1202

PART 1 - ATTORNEY/APPLICANT COPY

**ADENO-ASSOCIATED VECTOR COMPOSITIONS
FOR EXPRESSION OF FACTOR VIII**

COPY

RELATED APPLICATIONS

5 This application is a continuation of U.S. Application No. 09/470,618 filed December 22, 1999, which is a continuation-in-part of U.S. Application No. 09/364,862, filed July 30, 1999, which claims benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application Nos. 60/125,974 and 60/104,994, filed March 24, 1999 and October 20, 1998, respectively. All of these prior applications are hereby incorporated by reference in their
10 entireties.

FIELD OF THE INVENTION

 The present invention relates to AAV vectors suitable for hemophilia gene therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids encoding
15 Factor VIII into a recipient subject suffering from hemophilia A, such that the subject's blood is able to clot.

BACKGROUND

 Hemophilia is a genetic disease characterized by a blood clotting deficiency. In
20 hemophilia A (classic hemophilia, Factor VIII deficiency), an X-chromosome-linked genetic defect disrupts the gene encoding Factor VIII, a plasma glycoprotein, which is a key component in the blood clotting cascade. Human Factor VIII is synthesized as a single chain polypeptide, with a predicted molecular weight of 265 kDa. The Factor VIII gene codes for 2351 amino acids, and the protein has six domains, designated from the amino to
25 the carboxy terminus as A1-A2-B-A3-C1-C2 (Wood *et al.*, Nature 312:330 [1984]; Vehar *et al.*, Nature 312:337 [1984]; and Toole *et al.*, Nature 312:342 [1984]). Human Factor VIII is processed within the cell to yield a heterodimer primarily comprised of a heavy chain of 200 kDa containing the A1, A2, and B domains and an 80 kDa light chain containing the A3, C1, and C2 domains (Kaufman *et al.*, J. Biol. Chem., 263:6352-6362
30 [1988]). Both the single chain polypeptide and the heterodimer circulate in the plasma as inactive precursors (Ganz *et al.*, Eur. J. Biochem., 170:521-528 [1988]). Activation of

Factor VIII in plasma is initiated by thrombin cleavage between the A2 and B domains, which releases the B domain and results in a heavy chain consisting of the A1 and A2 domains. The 980 amino acid B domain is deleted in the activated procoagulant form of the protein. Additionally, in the native protein, two polypeptide chains ("a" and "b"),
5 flanking the B domain, are bound to a divalent calcium cation. Hemophilia may result from point mutations, deletions, or mutations resulting in a stop codon (*See, Antonarakis et al., Mol. Biol. Med., 4:81 [1987]*).

The disease is relatively rare, afflicting approximately one in 10,000 males. Hemophilia in females is extremely rare, although it may occur in female children of an affected father and carrier mother, as well as in females with X-chromosomal abnormalities (*e.g., Turner syndrome, X mosaicism, etc.*). The severity of each patient's disease is broadly characterized into three groups--"mild," "moderate," and "severe," depending on the severity of the patient's symptoms and circulating Factor VIII levels. While normal levels of Factor VIII range between 50 and 200 ng/mL plasma, mildly
15 affected patients have 6-60% of this value, and moderately affected patients have 1-5% of this value. Severely affected hemophiliacs have less than 1% of normal Factor VIII levels.

While hemophiliacs clearly require clotting factor after surgery or severe trauma, on a daily basis, spontaneous internal bleeding is a greater concern. Hemophiliacs experience spontaneous hemorrhages from early infancy, as well as frequent spontaneous
20 hemarthroses and other hemorrhages requiring clotting factor replacement. Without effective treatment, chronic hemophilic arthropathy occurs by young adulthood. Severely affected patients are prone to serious hemorrhages that may dissect through tissue planes, ultimately resulting in death due to compromised vital organs.

Hematomas are commonly observed in moderately and severely affected
25 hemophiliacs. In these patients, hematomas have a tendency to progressively enlarge and dissect in all directions. Some of these hematomas expand locally, resulting in local compression of adjacent organs, blood vessels, and nerves. A rare, yet often fatal, complication of abdominal hematomas is the perforation and drainage of the hematoma into the colon, resulting in infection and septicemia. Intracranial and/or extracranial
30 hemorrhage also represent very dangerous bleeding situations. While subcutaneous hematomas may dissect into muscle, pharyngeal and retropharyngeal hematomas (*e.g.,*

complicating bacterial or viral pharyngitis) may enlarge and obstruct the airway, sometimes resulting in a life-threatening situation that requires administration of a sufficient dose of Factor VIII concentrate to normalize the Factor VIII level.

5 In addition to hematomas, hemarthroses are commonly observed in hemophiliacs, with bleeding into the joint accounting for approximately 75% of hemophilic bleeding. Repeated hemorrhaging into the joints eventually results in extensive destruction of articular cartilage, synovial hyperplasia, and other reactive changes in adjacent tissues and bone. A major complication of repeated hemarthroses is joint deformity, which is often accompanied by muscle atrophy and soft tissue contractures; osteoporosis and cystic areas
10 in the subchondral bone may also develop, along with progressive loss of joint space.

Other symptoms are often observed in hemophiliacs, including hematuria and mucous membrane bleeding. Hematuria is experienced by virtually all severely affected hemophiliacs sometime during their lifetimes, and mucous membrane bleeding is common in hemophiliacs. Bone cysts (pseudotumors) are rare, but dangerous complications of
15 hemophilic bleeding. In many of these cases, immediate treatment is necessary.

In the early 1980s, many severely affected hemophiliacs were treated with Factor VIII concentrate about three times weekly. Unfortunately, these concentrates transmitted viruses, such as hepatitis B and/or C, and human immunodeficiency virus (HIV). In the United States and Western Europe, at least 75% of Factor VIII concentrate recipients have
20 been reported to have anti-HIV antibodies (*See, Schrier and Leung, supra*). Some of these patients also developed HIV-associated immune thrombocytopenia, a very serious complication in hemophiliacs. In spite of antiviral therapy (*e.g.*, with zidovudine and pentamidine prophylaxis), which has tended to slow disease progression, full-blown AIDS (acquired immunodeficiency syndrome) occurs at an inexorable rate in hemophiliacs
25 infected with HIV. Indeed, this has reversed the improvement in the life expectancy of hemophiliacs, which peaked at 66 years of age during the 1970s, and has dropped to 49 years (*See, Schrier and Leung, supra*). The development of virus-free preparations and recombinant Factor VIII has helped control infectious viral contamination.

30 However, for hemophiliacs, the availability of viral-free concentrates and recombinant Factor VIII, while significant, is but part of the solution. In order to prevent spontaneous internal bleeding episodes, patients suffering from hemophilia A must

consistently have serum Factor VIII levels of about 1%, and preferably 5%. Currently, the cost of viral-free concentrates and recombinant Factor VIII make it prohibitively expensive to administer the clotting factor prophylactically or on a maintenance basis. Indeed, most hemophiliacs in the U.S. do not receive recombinant Factor VIII therapy on a maintenance basis, but only receive it prior to activities or events which might cause bleeding (*e.g.*, surgery), or as a treatment for spontaneous bleeding.

Moreover, even if cost effective preparations of recombinant or virus-free Factor VIII were available, a steady state level of Factor VIII cannot be achieved by its daily administration. At best, patients receive widely varying levels of Factor VIII. Immediately following the administration, the levels are super-physiological, while prior to administration the levels are sub-physiological. Thus, there remains a need for methods and compositions that are relatively economic, yet effective in the treatment and prevention of bleeding in hemophiliacs, particularly spontaneous bleeds. Furthermore, there is a need in the art for methods and compositions for long term delivery of clotting factors (*e.g.*, Factor VIII) which more closely mimic the steady state physiological levels observed in normal individuals.

SUMMARY OF THE INVENTION

The present invention provides improved viral vectors suitable for gene therapy to treat hemophilia. In particular, the present invention provides AAV vectors and methods for treating hemophilia A by delivering nucleic acids coding for the clotting protein Factor VIII. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the vectors.

The present invention is particularly suited for use in hemophilia A gene therapy. Accordingly, in one embodiment of the invention, at least one AAV vector containing a nucleic acid molecule encoding Factor VIII is operably linked to control sequences that direct expression of Factor VIII in a suitable recipient cell. The AAV vectors are then introduced into a recipient cell of the subject, under conditions that result in expression of Factor VIII. The subject, therefore, has a continuous supply of Factor VIII available to clot blood during bleeding episodes.

Using the methods of the present invention, long term expression of therapeutic levels of Factor VIII have been achieved *in vivo*. In one embodiment, animals were administered, via the portal vein, two AAV vectors: one carrying the DNA sequence coding for the heavy chain of Factor VIII and the other carrying the DNA sequence coding for the light chain of Factor VIII. Blood samples were collected periodically and assayed for Factor VIII activity. Reproducibly, animals expressed between 600 and 900 ng/ml of biologically active Factor VIII, levels that are well above the normal physiological levels of approximately 200 ng/ml. Furthermore, these levels have been sustained for over 13 months without a decrease in Factor VIII levels or activity. In a related embodiment, a B-domain deleted form of Factor VIII was cloned into a single AAV vector and shown to express biologically active Factor VIII.

It is not intended, however, that the present invention be limited to specific embodiments. Many different forms of recombinant Factor VIII have been made and tested both *in vitro* and *in vivo*, using a variety of different control and regulatory sequences. Any DNA sequence coding for biologically active Factor VIII can be expressed using the AAV vectors and methods taught in the present invention. Therefore, the present invention encompasses any AAV vector or vectors containing Factor VIII sequences that produce biologically active Factor VIII protein *in vitro* or *in vivo*.

For example, in some embodiments, the AAV vector contains the first 57 base pairs of the Factor VIII heavy chain which encodes the 10 amino acid signal sequence, as well as the human growth hormone (hGH) polyadenylation sequence. In some alternative embodiments, the vector also contains the A1 and A2 domains, as well as 5 amino acids from the N-terminus of the B domain, and/or 85 amino acids of the C-terminus of the B domain, as well as the A3, C1, and C2 domains. In yet other embodiments, the nucleic acids coding for Factor VIII heavy chain and light chain were cloned into a single vector separated by 42 nucleic acids coding for 14 amino acids of the B domain.

The present invention also provides methods for administering the above-described vectors. For example, it is intended that the present invention encompass methods suitable for delivery of the AAV vectors to the livers of recipient patients or test animals. It is not intended that the present invention be limited to any particular route of administration.

However, in preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal or arterial vasculature.

These and other embodiments of the invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a schematic representation of the Factor VIII protein.

Figure 2 provides a schematic representation of a B-domain deleted form of Factor VIII protein.

10 Figure 3 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (AAV-F8-1) from internal terminal repeat (ITR to ITR), including control sequences.

15 Figure 4 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (PVM4.1c-F8AB) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 5 provides the sequence of pAAV-F8-1 (ITR to ITR), with the plasmid backbone omitted.

Figure 6 provides the sequence of pVm4.1cF8 Δ B (ITR to ITR), with the plasmid backbone omitted.

20 Figure 7 provides a map of rAAV-hFVIII-HC and rAAV-hFVIII-LC vectors.

Figure 8 provides a graph demonstrating the expression of various human FVIII constructs in mouse plasma.

Figure 9 provides Southern blot analyses of liver DNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

25 Figure 10 provides Southern blot analyses of DNA from different tissues using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

Figure 11 provides Northern blot analyses of liver RNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

GENERAL DESCRIPTION OF THE INVENTION

The present invention relates to improved viral vectors useful for expressing gene products at high levels in human cells. In particular, the present invention provides AAV vectors suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of Factor VIII protein in a host. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

The AAV vectors and rAAV virions of the present invention can be produced using standard methodology known to those of skill in the art. The methods generally involve the steps of: (1) introducing an AAV vector into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions. The AAV vector, AAV helper construct and the helper virus or accessory function vector(s) can be introduced into the host cell either simultaneously or serially, using standard transfection techniques.

Unless otherwise indicated, the practice of the present invention employs conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, including those described in such references as Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*; Glover (ed.) *DNA Cloning: A Practical Approach*, Vols. I and II; Gait (ed.) *Oligonucleotide Synthesis*; Hames and Higgins (eds.) *Nucleic Acid Hybridization*; Hames and Higgins (eds.) *Transcription and Translation*; Tijessen (ed.) *CRC Handbook of Parvoviruses*, Vols. I and II; and Fields and Knipe (eds.) *Fundamental Virology*, 2nd Edition, Vols. I and II.

Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

As used herein, the terms "gene transfer" and "gene delivery" refer to methods or systems for reliably inserting a particular nucleotide sequence (*e.g.*, DNA) into targeted cells. In particularly preferred embodiments, the nucleotide sequence comprises at least a portion of Factor VIII.

5 As used herein, the terms "vector," and "gene transfer vector" refer to any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus, virion, etc., which is capable of replication when associated with the proper control sequences and/or which can transfer nucleic acid sequences between cells. Thus, the term includes cloning and expression vectors, as well as viral vectors.

10 Gene transfer vectors may include transcription sequences such as polyadenylation sites, selectable markers or reporter genes, enhancer sequences, and other control sequences which allow for the induction of transcription. Such control sequences are described more fully below.

 The term "expression vector" as used herein refers to a recombinant DNA molecule
15 containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, as well as other sequences. Eukaryotic cells are generally known to utilize promoters (constitutive, inducible or tissue specific),
20 enhancers, and termination and polyadenylation signals, although some elements may be deleted and other elements added without sacrificing the necessary expression.

 As used herein, the terms "host" and "expression host" refer to organisms and/or cells which harbor an exogenous DNA sequence (*e.g.*, via transfection), an expression vector or vehicle, as well as organisms and/or cells that are suitable for use in expressing a
25 recombinant gene or protein. It is not intended that the present invention be limited to any particular type of cell or organism. Indeed, it is contemplated that any suitable organism and/or cell will find use in the present invention as a host.

 As used herein, the terms "viral replicons" and "viral origins of replication" refer to viral DNA sequences that allow for the extrachromosomal replication of a vector in a host
30 cell expressing the appropriate replication factors. In some embodiments, vectors which contain either the SV40 or polyoma virus origin of replication replicate to high copy

number, while vectors which contain the replicons from bovine papillomavirus or Epstein-Barr virus replicate extrachromosomally at low copy number may be utilized in other embodiments.

As used herein, the term "AAV vector" refers to a vector having functional or partly functional ITR sequences. The ITR sequences may be derived from an adeno-associated virus serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. The ITRs, however, need not be the wild-type nucleotide sequences, and may be altered (e.g., by the insertion, deletion or substitution of nucleotides), so long as the sequences retain function provide for functional rescue, replication and packaging. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or part, preferably the *rep* and/or *cap* genes but retain functional flanking ITR sequences. Functional ITR sequences are necessary for the rescue, replication and packaging of the AAV virion. Thus, an "AAV vector" is defined herein to include at least those sequences required in *cis* for replication and packaging (e.g., functional ITRs) of the virus.

As used herein, the term "ITR" refers to inverted terminal repeats. The terms "adeno-associated virus inverted terminal repeats" or "AAV ITRs" refer to the art-recognized palindromic regions found at each end of the AAV genome which function together in *cis* as origins of DNA replication and as packaging signals for the virus. For use in some embodiments of the present invention, flanking AAV ITRs are positioned 5' and 3' of one or more selected heterologous nucleotide sequences. Optionally, the ITRs together with the *rep* coding region or the Rep expression product provide for the integration of the selected sequences into the genome of a target cell.

As used herein, the term "AAV *rep* coding region" refers to the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. Muzyczka (Muzyczka, Curr. Top. Microbiol. Immunol., 158:97-129 [1992]) and Kotin (Kotin, Hum. Gene Ther., 5:793-801 [1994]) provide additional descriptions of the

AAV *rep* coding region, as well as the *cap* coding region described below. Suitable homologues of the AAV *rep* coding region include the human herpesvirus 6 (HHV-6) *rep* gene which is also known to mediate AAV-2 DNA replication (Thomson *et al.*, Virol., 204:304-311 [1994]).

5 As used herein, the term "AAV *cap* coding region" refers to the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These *cap* expression products supply the packaging functions which are collectively required for packaging the viral genome.

10 As used herein, the term "AAV helper function" refers to AAV coding regions capable of being expressed in the host cell to complement AAV viral functions missing from the AAV vector. Typically, the AAV helper functions include the AAV *rep* coding region and the AAV *cap* coding region. An "AAV helper construct" is a vector containing AAV coding regions required to complement AAV viral functions missing from the AAV vector (*e.g.*, the AAV *rep* coding region and the AAV *cap* coding region).

15 As used herein, the terms "accessory functions" and "accessory factors" refer to functions and factors that are required by AAV for replication, but are not provided by the AAV virion (or rAAV virion) itself. Thus, these accessory functions and factors must be provided by the host cell, a virus (*e.g.*, adenovirus or herpes simplex virus), or another expression vector that is co-expressed in the same cell. Generally, the E1, E2A, E4 and
20 VA coding regions of adenovirus are used to supply the necessary accessory function required for AAV replication and packaging (Matsushita *et al.*, Gene Therapy 5:938 [1998]).

25 As used herein, the term "wild type" ("wt") refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that
30 naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "AAV virion" refers to a complete virus particle, such as a "wild-type" (wt) AAV virus particle (comprising a linear, single-stranded AAV nucleic acid genome associated with an AAV capsid protein coat). In this regard, single-stranded AAV nucleic acid molecules of either complementary sense (e.g., "sense" or "antisense" strands), can be packaged into any one AAV virion and both strands are equally infectious.

As used herein, the terms "recombinant AAV virion," and "rAAV virion" refer to an infectious viral particle containing a heterologous DNA molecule of interest (e.g., Factor VIII sequence) which is flanked on both sides by AAV ITRs. In some embodiments of the present invention, an rAAV virion is produced in a suitable host cell which contains an AAV vector, AAV helper functions and accessory functions introduced therein. In this manner, the host cell is rendered capable of encoding AAV polypeptides that are required for packaging the AAV vector containing a recombinant nucleotide sequence of interest, such as at least a portion of Factor VIII or portions of Factor VIII domains, into recombinant virion particles for subsequent gene delivery.

As used herein, the term "transfection" refers to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art (See e.g., Graham *et al.*, *Viol.*, 52:456 [1973]; Sambrook *et al.*, *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York [1989]; Davis *et al.*, *Basic Methods in Molecular Biology*, Elsevier, [1986]; and Chu *et al.*, *Gene* 13:197 [1981]. Such techniques can be used to introduce one or more exogenous DNA moieties, such as a gene transfer vector and other nucleic acid molecules, into suitable recipient cells.

As used herein, the terms "stable transfection" and "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA.

As used herein, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the

regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

5 As used herein, the term "transduction" denotes the delivery of a DNA molecule to a recipient cell either *in vivo* or *in vitro*, via a replication-defective viral vector, such as via a recombinant AAV virion.

10 As used herein, the term "recipient cell" refers to a cell which has been transfected or transduced, or is capable of being transfected or transduced, by a nucleic acid construct or vector bearing a selected nucleotide sequence of interest (*i.e.*, Factor VIII). The term includes the progeny of the parent cell, whether or not the progeny are identical in morphology or in genetic make-up to the original parent, so long as the selected nucleotide sequence is present.

15 The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence
20 in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, synthetic sequences having codons different from the native gene). Similarly, a cell transfected with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to
25 heterologous DNA, as used herein.

As used herein, "coding sequence" or a sequence which "encodes" a particular antigen, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo*, when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are
30 determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from

prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

As used herein, the term "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

As used herein, the term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

As used herein, the term "regulatory element" refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

The term DNA "control sequences" refers collectively to regulatory elements such as promoter sequences, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need

always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate recipient cell.

Transcriptional control signals in eukaryotes generally comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (Maniatis *et al.*, Science 236:1237 [1987]). Promoter and enhancer elements have been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control sequences, *i.e.*, promoters, are also found in prokaryotes). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest (*i.e.*, Factor VIII). Some eukaryotic promoters and enhancers have a broad host range while others are functional in a limited subset of cell types (*See e.g.*, Voss *et al.*, Trends Biochem. Sci., 11:287 [1986]; and Maniatis *et al.*, *supra*, for reviews). For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (Dijkema *et al.*, EMBO J. 4:761 [1985]). Two other examples of promoter and enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 α gene (Uetsuki *et al.*, J. Biol. Chem., 264:5791 [1989]; Kim *et al.*, *Gene* 91:217 [1990]; and Mizushima and Nagata, Nucl. Acids. Res., 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777 [1982]) and the human cytomegalovirus (Boshart *et al.*, Cell 41:521 [1985]). Promoters and enhancers can be found naturally alone or together. For example, the long terminal repeats of retroviruses contain both promoter and enhancer functions. Moreover, generally promoters and enhancers act independently of the gene being transcribed or translated. Thus, the enhancer and promoter may be "endogenous" or "exogenous" or "heterologous." An "endogenous" enhancer/promoter is one which is naturally linked with a given gene in the genome. An "exogenous" or "heterologous" enhancer and promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter.

As used herein, the term "tissue specific" refers to regulatory elements or control sequences, such as a promoter, enhancers, etc., wherein the expression of the nucleic acid sequence is substantially greater in a specific cell type(s) or tissue(s). In particularly preferred embodiments, the albumin promoter and the transthyretin promoter display increased expression of FVIII in hepatocytes, as compared to other cell types. It is not intended, however, that the present invention be limited to the albumin or transthyretin promoters or to hepatic-specific expression, as other tissue specific regulatory elements, or regulatory elements that display altered gene expression patterns, are contemplated.

The presence of "splicing signals" on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells requires expression of signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site" or "poly A sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A signal is contained on a 237 bp *Bam*HI/*Bcl*II restriction fragment and directs both termination and polyadenylation (Sambrook *et al.*, *supra*, at 16.6-16.7).

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding

sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids are nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. The isolated nucleic acid, oligonucleotide, or polynucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid, oligonucleotide or polynucleotide is to be utilized to express a protein, the oligonucleotide or polynucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide or polynucleotide may single-stranded), but may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide or polynucleotide may be double-stranded).

As used herein, the term "purified" or "to purify" refers to the removal of contaminants from a sample. For example, antibodies may be purified by removal of contaminating non-immunoglobulin proteins; they may also purified by the removal of immunoglobulin that does not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII). The removal of non-immunoglobulin proteins and/or the removal of immunoglobulins that do not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII) results in an increase in the percent of desired antigen-reactive immunoglobulins in the sample. In another example, recombinant polypeptides of Factor VIII are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant polypeptides is thereby increased in the sample.

As used herein, the term "chimeric protein" refers to two or more coding sequences obtained from different genes, that have been cloned together and that, after translation, act as a single polypeptide sequence. Chimeric proteins are also referred to as "hybrid proteins." As used herein, the term "chimeric protein" refers to coding sequences that are obtained from different species of organisms, as well as coding sequences that are obtained from the same species of organisms.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The composition may comprise an aqueous solution.

As used herein, the term "at risk" is used in references to individuals who are at risk for experiencing hemorrhagic episodes. In particularly preferred embodiments, the individuals are hemophiliacs with mild, moderate, or severe hemophilia.

As used herein, the term "subject" refers to any animal (*i.e.*, vertebrates and invertebrates), while the term "vertebrate subject" refers to any member of the subphylum Chordata. It is intended that the term encompass any member of this subphylum, including, but not limited to humans and other primates, rodents (*e.g.*, mice, rats, and guinea pigs), lagamorphs (*e.g.*, rabbits), bovines (*e.g.*, cattle), ovines (*e.g.*, sheep), caprines (*e.g.*, goats), porcines (*e.g.*, swine), equines (*e.g.*, horses), canines (*e.g.*, dogs), felines (*e.g.*, cats), domestic fowl (*e.g.*, chickens, turkeys, ducks, geese, other gallinaceous birds, etc.), as well as feral or wild animals, including, but not limited to, such animals as ungulates (*e.g.*, deer), bear, fish, lagamorphs, rodents, birds, etc. It is not intended that the term be limited to a particular age or sex. Thus, adult and newborn subjects, as well as fetuses, whether male or female, are encompassed by the term.

As defined herein, a "therapeutically effective amount" or "therapeutic effective dose" is an amount or dose of AAV vector or virions capable of producing sufficient amounts of Factor VIII to decrease the time it takes for a subject's blood to clot. Generally, severe hemophiliacs having less than 1% of normal levels of FVIII have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs.

DETAILED DESCRIPTION OF THE INVENTION

Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an AAV vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended.

A. Control Sequences

In some embodiments of the present invention, heterologous control sequences are employed with the vectors. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), a herpes simplex virus (HSV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE), a rous sarcoma virus (RSV) promoter, synthetic promoters, hybrid promoters, and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, also find use herein. Such promoter sequences are commercially available (*e.g.*, from Stratagene).

It is contemplated that in some embodiments, tissue-specific expression may be desirable (*e.g.*, expression of biologically active Factor VIII by hepatocytes). It is not intended that the present invention be limited to expression of biologically active Factor VIII by any particular cells or cell types. However, as hepatocytes (*i.e.*, liver cells) are the cells that normally synthesized Factor VIII (*See*, Kaufman, *Ann. Rev. Med.*, 43:325 [1992]), it is contemplated that in some particularly preferred embodiments, the compositions of the present invention be administered to the liver.

In preferred embodiments, expression is achieved by coupling the coding sequence for Factor VIII with heterologous control sequences derived from genes that are specifically transcribed by a selected tissue type. A number of tissue-specific promoters have been described above which enable directed expression in selected tissue types. However, control sequences used in the present AAV vectors can also comprise control sequences normally associated with the selected nucleic acid sequences.

B. Construction of AAV Factor VIII Vectors

AAV vectors that contain a control sequence and a nucleotide sequence of interest (*i.e.*, at least a portion of the sequence encoding Factor VIII), bounded by AAV ITRs (*i.e.*, AAV vectors), can be constructed by directly inserting selected sequences into an AAV genome with the major AAV open reading frames ("ORFs") excised. Other portions of the AAV genome can also be deleted, so long as a sufficient portion of the ITRs remain to allow for replication and packaging functions. These constructs can be designed using techniques well known in the art (*See e.g.*, U.S. Patent Nos. 5,173,414 and 5,139,941, all of which are herein incorporated by reference); International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski *et al.*, *Mol. Cell. Biol.*, 8:3988-3996 [1988]; Vincent *et al.*, *Vaccines 90* [Cold Spring Harbor Laboratory Press, 1990]; Carter, *Curr. Opin. Biotechnol.*, 3:533-539 [1992]; Muzyczka, *Curr. Top. Microbiol. Immunol.*, 158:97-129 [1992]; Kotin, *Hum. Gene Ther.*, 5:793-801 [1994]; Shelling and Smith, *Gene Ther.*, 1:165-169 [1994]; and Zhou *et al.*, *J. Exp. Med.*, 179:1867-1875 [1994]).

Alternatively, AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct that is present in another vector using standard ligation techniques, such as those described in Sambrook *et al.*, *supra*. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration). AAV vectors which contain ITRs have been described in (*e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference). In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC") under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

Additionally, chimeric genes can be produced synthetically to include AAV ITR sequences arranged 5' and 3' of a selected nucleic acid sequence. The complete chimeric sequence is assembled from overlapping oligonucleotides prepared by standard methods

(See e.g., Edge, Nature 292:756 [1981]; Nambair *et al.*, Science 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Moreover, it is not intended that the present invention be limited to any specific Factor VIII sequence. Many natural and recombinant forms of Factor VIII have been isolated and assayed both *in vitro* and *in vivo*, using a variety of different regulatory elements and control sequences. Therefore, any known, or later discovered, DNA sequence coding for biologically active Factor VIII can be expressed, alone or in combination with at least one additional vector, using the AAV vectors and methods taught in the present invention. Examples of naturally occurring and recombinant forms of Factor VIII can be found in the patent and scientific literature including, U.S. 5,563,045, U.S. 5,451,521, U.S. 5,422,260, U.S. 5,004,803, U.S. 4,757,006, U.S. 5,661,008, U.S. 5,789,203, U.S. 5,681,746, U.S. 5,595,886, U.S. 5,045,455, U.S. 5,668,108, U.S. 5,633,150, U.S. 5,693,499, U.S. 5,587,310, U.S. 5,171,844, U.S. 5,149,637, U.S. 5,112,950, U.S. 4,886,876, WO 94/11503, WO 87/07144, WO 92/16557, WO 91/09122, WO 97/03195, WO 96/21035, WO 91/07490, EP 0 672 138, EP 0 270 618, EP 0 182 448, EP 0 162 067, EP 0 786 474, EP 0 533 862, EP 0 506 757, EP 0 874 057, EP 0 795 021, EP 0 670 332, EP 0 500 734, EP 0 232 112, EP 0 160 457, Sanberg *et al.*, XXth Int. Congress of the World Fed. Of Hemophilia (1992), and Lind *et al.*, Eur. J. Biochem., 232:19 (1995).

Nucleic acid sequences coding for the above-described Factor VIII can be obtained using recombinant methods, such as by screening cDNA and genomic libraries from cells expressing Factor VIII or by deriving the sequence from a vector known to include the same. Furthermore, the desired sequence can be isolated directly from cells and tissues containing the same, using standard techniques, such as phenol extraction and PCR of cDNA or genomic DNA (See e.g., Sambrook *et al.*, *supra*, for a description of techniques used to obtain and isolate DNA). Nucleotide sequences encoding an antigen of interest (*i.e.*, Factor VIII sequence) can also be produced synthetically, rather than cloned. The complete sequence can be assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence (See e.g., Edge, Nature 292:756 [1981]; Nambair *et al.*, Science 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Although it is not intended that the present invention be limited to any particular methods for assessing the production of biologically active Factor VIII, such methods as immunoassays (*e.g.*, ELISA) and biological activity assays are contemplated (*e.g.*, coagulation activity assays).

5 Furthermore, while in particularly preferred embodiments, human Factor VIII is encompassed by the present invention, it is not intended that the present invention be limited to human Factor VIII. Indeed, it is intended that the present invention encompass Factor VIII from animals other than humans, including but not limited to companion animals (*e.g.*, canines, felines, and equines), livestock (*e.g.*, bovines, caprines, and ovines),
10 laboratory animals (*e.g.*, rodents such as murines, as well as lagamorphs), and "exotic" animals (*e.g.*, marine mammals, large cats, etc.).

II. Virion Production

Producing AAV Factor VIII vectors and rAAV Factor VIII virions of the present
15 invention generally involve the steps of: (1) introducing an AAV vector containing the Factor VIII gene into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell,
20 wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions.

The above-described vectors and constructs can be introduced into a cell using standard methodology known to those of skill in the art (*e.g.*, transfection). A number of
25 transfection techniques are generally known in the art (*See e.g.*, Graham *et al.*, Virol., 52:456 [1973], Sambrook *et al. supra*, Davis *et al., supra*, and Chu *et al.*, Gene 13:197 [1981]). Particularly suitable transfection methods include calcium phosphate co-precipitation (Graham *et al.*, Virol., 52:456-467 [1973]), direct micro-injection into cultured cells (Capecchi, Cell 22:479-488 [1980]), electroporation (Shigekawa *et al.*,
30 BioTechn., 6:742-751 [1988]), liposome-mediated gene transfer (Mannino *et al.*, BioTechn., 6:682-690 [1988]), lipid-mediated transduction (Felgner *et al.*, Proc. Natl.

Acad. Sci. USA 84:7413-7417 [1987]), and nucleic acid delivery using high-velocity microprojectiles (Klein *et al.*, Nature 327:70-73 [1987]).

For the purposes of the invention, suitable host cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule. The term includes the progeny of the original cell which has been transfected. Thus, as indicated above, a "host cell" as used herein generally refers to a cell which has been transfected with an exogenous DNA sequence. Cells from the stable human cell line, 293 (ATCC Accession No. CRL1573) are preferred in the practice of the present invention. Particularly, the human cell line 293 is a human embryonic kidney cell line that has been transformed with adenovirus type-5 DNA fragments (Graham *et al.*, J. Gen. Virol., 36:59 [1977]), and expresses the adenoviral E1a and E1b genes (Aiello *et al.*, Virol., 94:460 [1979]). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

Host cells containing the above-described AAV vectors must be rendered capable of providing AAV helper functions in order to replicate and encapsidate the nucleotide sequences flanked by the AAV ITRs to produce rAAV virions. AAV helper functions are generally AAV-derived coding sequences which can be expressed to provide AAV gene products that, in turn, function in *trans* for productive AAV replication. AAV helper functions are used herein to complement necessary AAV functions that are missing from the AAV vectors. Thus, AAV helper functions include one, or both of the major AAV ORFs, namely the *rep* and *cap* coding regions, or functional homologues thereof.

AAV helper functions are introduced into the host cell by transfecting the host cell with an AAV helper construct either prior to, or concurrently with, the transfection of the AAV vector. AAV helper constructs are thus used to provide at least transient expression of AAV *rep* and/or *cap* genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

In preferred embodiments, these constructs are in the form of a vector, including, but not limited to, plasmids, phages, transposons, cosmids, viruses, or virions. A number of AAV helper constructs have been described, such as the commonly used plasmids

pAAV/Ad and pIM29+45 which encode both Rep and Cap expression products (*See e.g.*, Samulski *et al.*, J. Virol., 63:3822-3828 [1989]; and McCarty *et al.*, J. Virol., 65:2936-2945 [1991]). A number of other vectors have been described which encode Rep and/or Cap expression products (*See e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference).

Both AAV vectors and AAV helper constructs can be constructed to contain one or more optional selectable markers. Suitable markers include genes which confer antibiotic resistance or sensitivity to, impart color to, or change the antigenic characteristics of those cells which have been transfected with a nucleic acid construct containing the selectable marker when the cells are grown in an appropriate selective medium. Several selectable marker genes that are useful in the practice of the invention include the gene encoding aminoglycoside phosphotransferase (APH) that allows selection in mammalian cells by conferring resistance to G418 (Sigma). Other suitable markers are known to those of skill in the art.

The host cell (or packaging cell) must also be rendered capable of providing non-AAV derived functions, or "accessory functions," in order to produce rAAV virions. Accessory functions are non-AAV derived viral and/or cellular functions upon which AAV is dependent for its replication. Thus, accessory functions include at least those non-AAV proteins and RNAs that are required in AAV replication, including those involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of *rep* and *cap* expression products and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses.

Particularly, accessory functions can be introduced into and then expressed in host cells using methods known to those of skill in the art. Commonly, accessory functions are provided by infection of the host cells with an unrelated helper virus. A number of suitable helper viruses are known, including adenoviruses; herpesviruses such as herpes simplex virus types 1 and 2; and vaccinia viruses. Nonviral accessory functions will also find use herein, such as those provided by cell synchronization using any of various known agents (*See e.g.*, Buller *et al.*, J. Virol., 40:241-247 [1981]; McPherson *et al.*, Virol., 147:217-222 [1985]; and Schlehofer *et al.*, Virol., 152:110-117 [1986]).

Alternatively, accessory functions can be provided using an accessory function vector. Accessory function vectors include nucleotide sequences that provide one or more accessory functions. An accessory function vector is capable of being introduced into a suitable host cell in order to support efficient AAV virion production in the host cell.

5 Accessory function vectors can be in the form of a plasmid, phage, virus, transposon or cosmid. Accessory vectors can also be in the form of one or more linearized DNA or RNA fragments which, when associated with the appropriate control sequences and enzymes, can be transcribed or expressed in a host cell to provide accessory functions.

10 Nucleic acid sequences providing the accessory functions can be obtained from natural sources, such as from the genome of adenovirus, or constructed using recombinant or synthetic methods known in the art. In this regard, adenovirus-derived accessory functions have been widely studied, and a number of adenovirus genes involved in accessory functions have been identified and partially characterized (*See e.g.*, Carter, "Adeno-Associated Virus Helper Functions," in *CRC Handbook of Parvoviruses*, Vol. I (P.

15 Tijssen, ed.) [1990], and Muzyczka, *Curr. Top. Microbiol. Immun.*, 158:97-129 [1992]). Specifically, early adenoviral gene regions E1a, E2a, E4, VAI RNA and, possibly, E1b are thought to participate in the accessory process (Janik *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1925-1929 [1981]). Herpesvirus-derived accessory functions have been described (*See e.g.*, Young *et al.*, *Prog. Med. Virol.*, 25:113 [1979]). Vaccinia virus-derived accessory

20 functions have also been described (*See e.g.*, Carter, *supra.*, and Schlehofer *et al.*, *Virol.*, 152:110-117 [1986]).

As a consequence of the infection of the host cell with a helper virus, or transfection of the host cell with an accessory function vector, accessory functions are expressed which transactivate the AAV helper construct to produce AAV Rep and/or Cap

25 proteins. The Rep expression products direct excision of the recombinant DNA (including the DNA of interest encoding at least a portion of Factor VIII) from the AAV vector. The Rep proteins also serve to duplicate the AAV genome. The expressed Cap proteins assemble into capsids, and the recombinant AAV genome is packaged into the capsids. Thus, productive AAV replication ensues, and the DNA is packaged into rAAV virions.

30 Following recombinant AAV replication, rAAV virions can be purified from the host cell using a variety of conventional purification methods, such as CsCl gradients.

Further, if helper virus infection is employed to express the accessory functions, residual helper virus can be inactivated, using known methods. For example, adenovirus can be inactivated by heating to temperatures of approximately 60°C for approximately 20 minutes or more, as appropriate. This treatment selectively inactivates the helper adenovirus which is heat labile, while preserving the rAAV which is heat stable.

III. Pharmaceutical Compositions

The resulting rAAV virions are then ready for use in pharmaceutical compositions which can be delivered to a subject, so as to allow production of biologically active Factor VIII. Pharmaceutical compositions comprise sufficient genetic material that allows the recipient to produce a therapeutically effective amount of Factor VIII so as to reduce, stop and/or prevent hemorrhage. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, clotting factors or factor precursors, drugs or hormones. In preferred embodiments, the pharmaceutical compositions also contain a pharmaceutically acceptable excipient. Such excipients include any pharmaceutical agent that does not itself induce an immune response harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol, sugars and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., N.J. [1991]).

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as

Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art (*e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes).

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment. For administration of Factor VIII-containing vectors, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. Determining a therapeutic effective dose is well within the capability of those skilled in the art using the techniques taught in the present invention, such as ELISA and ChromZ FVIII coagulation activity assay, and other techniques known in the art.

Therapeutic doses will depend on, among other factors, the age and general condition of the subject, the severity of hemophilia, and the strength of the control sequences. Thus, a therapeutically effective amount in humans will fall in a relatively broad range that can be determined through clinical trials.

5 It is intended that the dosage treatment and regimen used with the present invention will vary, depending upon the subject and the preparation to be used. Thus, the dosage treatment may be a single dose schedule or a multiple dose schedule. Moreover, the subject may be administered as many doses as appropriate to achieve or maintain the desired blood clotting time.

10 Direct delivery of the pharmaceutical compositions *in vivo* will generally be accomplished via injection using a conventional syringe, although other delivery methods such as convention-enhanced delivery are envisioned (*See e.g.*, United States Patent No. 5,720,720, incorporated herein by reference). In this regard, the compositions can be delivered subcutaneously, epidermally, intradermally, intrathecally, intraorbitally,
15 intramucosally (*e.g.*, nasally, rectally and vaginally), intraperitoneally, intravenously, intraarterially, orally, or intramuscularly. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. In particularly preferred embodiments, the compositions are administered intravenously in the portal vasculature or hepatic artery

20 One skilled in the art will recognize that the methods and compositions described above are also applicable to a range of other treatment regimens known in the art. For example, the methods and compositions of the present invention are compatible with *ex vivo* therapy (*e.g.*, where cells are removed from the body, incubated with the AAV vector and the treated cells are returned to the body).

25 IV. ADMINISTRATION

AAV vector can be administered to any tissue suitable for the expression of Factor VIII. In a preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal vasculature or hepatic artery where it is thought,
30 without being bound by theory, that the vector transduces hepatocytes. Current approaches to targeting genes to the liver have focused upon *ex vivo* gene therapy. *Ex vivo*

liver-directed gene therapy involves the surgical removal of liver cells, transduction of the liver cells *in vitro* (e.g., infection of the explanted cells with recombinant retroviral vectors) followed by injection of the genetically modified liver cells into the liver or spleen of the patient. A serious drawback for *ex vivo* gene therapy of the liver is the fact that hepatocytes cannot be maintained and expanded in culture. Therefore, the success of *ex vivo* liver-directed gene therapy depends upon the ability to efficiently and stably engraft the genetically modified (*i.e.*, transduced) hepatocytes and their progeny. It has been reported that even under optimal conditions, autologous modified liver cells injected into the liver or spleen which engraft represent only a small percentage (less than 10%) of the total number of cells in the liver. Ectopic engraftment of transduced primary hepatocytes into the peritoneal cavity has been tried, in order to address the problem of engraftment in the liver.

Given the problems associated with *ex vivo* liver-directed gene therapy, *in vivo* approaches have been investigated for the transfer of genes into hepatocytes, including the use of recombinant retroviruses, recombinant adenoviruses, liposomes and molecular conjugates. While these *in vivo* approaches do not suffer from the drawbacks associated with *ex vivo* liver-directed gene therapy, they do not provide a means to specifically target hepatocytes. In addition, several of these approaches require performance of a partial hepatectomy, in order to achieve prolonged expression of the transferred genes *in vivo*. Adenovirus and molecular conjugate based delivery methods also result in liver toxicity and inflammation which is an undesirable feature of Factor VIII gene therapy. The present invention provides compositions and methods for the long-term expression of biologically active Factor VIII. It is contemplated that the present invention will bypass the need for partial hepatectomy, while allowing expression of Factor VIII in concentrations that are therapeutic *in vivo*. The present invention further provides gene therapy compositions and methods that target hepatocytes for the production of Factor VIII by treated individuals.

Other tissues, however, may be suitable for the expression of Factor VIII even if they are not the tissue that normally synthesizes the protein. Muscle cells, for example, have been shown to express biologically active blood clotting Factor IX even though it is normally synthesized in the liver.

Finally, the AAV vectors may contain any nucleic acid sequences coding for biologically active Factor VIII. Additionally, the AAV vectors may contain a nucleic acid coding for fragments of Factor VIII which is itself not biologically active, yet when administered into the subject improves or restores the blood clotting time. For example, as discussed above, the Factor VIII protein comprises two polypeptide chains: a heavy chain and a light chain separated by a B-domain which is cleaved during processing. As demonstrated by the present invention, co-transducing recipient cells with the Factor VIII heavy and light chains leads to the expression of biologically active Factor VIII. Because, however, most hemophiliacs contain a mutation or deletion in only one of the chains (*e.g.*, heavy or light chain), it may be possible to administer only the chain defective in the patient and allow the patient to supply the other chain. In this case, the AAV vector would fall within the scope of the invention even though the single chain (*i.e.*, heavy or light) would not be biologically active until it was administered into a subject which can supply the second chain, thus forming biologically active Factor VIII.

V. FACTOR VIII ASSAYS

As described in the Experimental section below, there are many ways to assay Factor VIII expression and activity. Although the present invention is not limited to immunoassay methods, the present invention also provides methods for detecting Factor VIII expression comprising the steps of: a) providing a sample suspected of containing Factor VIII, and a control containing a known amount of known Factor VIII; and b) comparing the test sample with the known control, to determine the relative concentration of Factor VIII in the sample. Thus, the methods are capable of identifying samples (*e.g.*, patient samples) with sufficient or insufficient quantities of Factor VIII. In addition, the methods may be conducted using any suitable means to determine the relative concentration of Factor VIII in the test and control samples, including but not limited to means selected from the group consisting of Western blot analysis, Northern blot analysis, Southern blot analysis, denaturing polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), reverse transcriptase-coupled polymerase chain reaction (RT-PCR), enzyme-linked

immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescent immunoassay (IFA). Thus, the methods may be conducted to determine the presence of normal Factor VIII sequences in the genome of the animal source of the test sample, or the expression of Factor VIII (mRNA or protein), as well as detect the presence of abnormal or mutated
5 Factor VIII gene sequences in the test samples.

In one preferred embodiment, the presence of Factor VIII is detected by immunochemical analysis. For example, the immunochemical analysis can comprise detecting binding of an antibody specific for an epitope of Factor VIII. In one another preferred embodiment of the method, the antibody comprises polyclonal antibodies, while
10 in another preferred embodiment, the antibody comprises monoclonal antibodies.

It is further contemplated that antibodies directed against at least a portion of Factor VIII will be used in methods known in the art relating to the localization and structure of Factor VIII (*e.g.*, for Western blotting), measuring levels thereof in appropriate biological samples, etc. The antibodies can be used to detect Factor VIII in a biological sample from
15 an individual (*e.g.*, an individual treated using the methods and/or compositions of the present invention). The biological sample can be a biological fluid, including, but not limited to, blood, serum, plasma, interstitial fluid, urine, cerebrospinal fluid, synovial fluid, and the like. In particular, the antigen can be detected from cellular sources, including, but not limited to, hepatocytes. For example, cells can be obtained from an individual and
20 lysed (*e.g.*, by freeze-thaw cycling, or treatment with a mild cytolytic detergent including, but not limited to, TRITON X-100, digitonin, NONIDET P (NP)-40, saponin, and the like, or combinations thereof; *See, e.g.*, International Patent Publication WO 92/08981).

The biological samples can then be tested directly for the presence of the Factor VIII using an appropriate strategy (*e.g.*, ELISA or RIA) and format (*e.g.*, microwells,
25 dipstick [*e.g.*, as described in International Patent Publication WO 93/03367], etc.). Alternatively, proteins in the sample can be size separated (*e.g.*, by polyacrylamide gel electrophoresis (PAGE), with or without sodium dodecyl sulfate (SDS), and the presence of Factor VIII detected by immunoblotting [*e.g.*, Western blotting]). Immunoblotting techniques are generally more effective with antibodies generated against a peptide
30 corresponding to an epitope of a protein, and hence, are particularly suited to the present invention. In another preferred embodiment, the level of Factor VIII is assayed using the

whole-blood clotting time and activated partial thromboplastin time (aPTT) of the subject's blood using techniques well known in the art (Herzog *et al.*, Nature Medicine 5:56 [1999]).

5 The foregoing explanations of particular assay systems are presented herein for purposes of illustration only, in fulfillment of the duty to present an enabling disclosure of the invention. It is to be understood that the present invention contemplates a variety of immunochemical assay protocols within its spirit and scope. Indeed, other methods such as biological assays to determine the presence and activity of Factor VIII are also encompassed by the present invention.

10 Thus, in addition to the immunoassay systems described above, other assay systems, such as those designed to measure and/or detect Fraction VIII and/or clotting ability of a subject's blood are also encompassed by the present invention (*e.g.*, the ChromZ FVIII coagulation activity [FVIII-c] assay [Helena Labs]).

EXPERIMENTAL

15 Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

20 Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

In the experimental disclosure which follows, the following abbreviations apply: N (Normal); M (Molar); mM (millimolar); μ M (micromolar); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); mU (milliunits);
25 51 Cr (Chromium 51); μ Ci (microcurie); EC (degrees Centigrade); hFVIII (human factor VIII); FVIII (factor VIII); pH (hydrogen ion concentration); JRH grade; NaCl (sodium chloride); HCl (hydrochloric acid); OD (optical density); bp (base pair(s)); ATP (adenosine 5'-triphosphate); PCR (polymerase chain reaction); DNA (deoxyribonucleic acid); cDNA (complementary DNA); AAV (adeno-associated virus); rAAV (recombinant adeno-associated virus);
30 ITR (inverted terminal repeat); FCS or FBS (fetal calf serum; fetal

bovine serum); CFA (complete Freund's adjuvant); BSA (bovine serum albumin); ATCC (American Type Culture Collection, Rockville, MD); Sigma (Sigma Aldrich, St. Louis, MO); Biodesign International (Biodesign International, Kennebunk, MI); Baxter Hyland (Baxter Healthcare Corp., Biotech Group--Hyland Division, Hayward, CA); Helena Labs (Helena Laboratories, Beaumont, TX); American Diagnostica (American Diagnostica, Greenwich, CT); Accurate Chemical (Accurate Chemical and Scientific Corp., Westbury, NY); Molecular Probes (Molecular Probes, Eugene, OR); Vysis (Vysis, Downer Grove, IL); Tel-Test (Tel-Test, Inc., Friendswood, TX); Molecular Dynamics (Molecular Dynamics, Sunnyvale, CA); NUNC (Naperville, IL); and Stratagene (Stratagene Cloning Systems, La Jolla, CA); Affinity Biologicals (Affinity Biologicals, Inc., Hamilton, Ontario); and Biodesign (Biodesign International, Kennebunkport, ME).

EXAMPLE 1

Dual Vector Plasmid Construction

The heavy and light chains of human Factor VIII (hFVIII) were assembled according to those reported by Yonemura *et al* (Yonemura *et al.*, Prot. Engineer., 6:669-674 [1993]) and cloned as expression cassettes into AAV vectors. Both vectors contain the promoter and the first non-coding intron (from -573 to +985) from the human elongation factor 1 α (EF1 α) gene (Uetsuki *et al*, J. Biol. Chem., 264:5791-5798 [1989]; and Kim *et al.*, Gene 9:217-223 [1990]). Each vector also contains the first 57 base pairs of the FVIII heavy chain encoding the 19 amino acid signal sequence. The heavy chain construct encodes the A1 and A2 domains and 5 amino acids from the N terminus of the B domain. The light chain vector encodes 85 amino acids of the carboxy terminal B domain, in addition to the A3, C1, and C2 domains. Both vectors utilize the human growth hormone (hGH) polyadenylation signal. The expression cassettes were inserted between AAV ITRs. The initial cloning step involved deleting 854 bp of EF1 α sequences between the *SpeI* and *XcmI* sites of pVm4.1e-hFIX (Nakai *et al.*, Blood 91:1-9 [1998]), and religating to create pVm4.1e δ D-hFIX.

This construct was then digested with *EcoRI*, which released the hFIX cDNA, and was ligated to an oligonucleotide containing *MfeI* ends (*EcoRI*-compatible) and an internal

*Cla*I restriction site, creating pVm4.1e δ D-linker. The heavy and light chain fragments, including the hGH polyadenylation sequences were isolated from pVm4.1cFVIII-HC and pVm4.1cFVIII-LC, respectively as *Cla*I-*Bst*EII fragments. These fragments were cloned between the corresponding sites in the pVm4.1e δ D-linker, creating plasmids pVm4.1e δ D-FVIII-HC (also, rAAV-hFVIII-HC) and pVm4.1e δ D-FVIII-LC (also, rAAV-hFVIII-LC).

Figure 7 provides a map of the constructs. In this figure, the upper line in each panel represents the gene structure of the vectors, and the lower line represents the structure of the hFVIII protein domains encoded by the vectors (ITR, AAV inverted terminal repeat; EF1 α Pro/Intron 1, human polypeptide elongation factor 1 α gene promoter and first intron; hFVIII-HC human FVIII cDNA; hFVIII-LC, human FVIII cDNA; hGH PA, human growth hormone polyadenylation signal; SS, human FVIII signal sequence; A1, A2, "B", A3, C1, C2, complete and incomplete (") protein domains of the hFVIII protein).

EXAMPLE 2

Single Vector Plasmid Construction

The plasmid pAAV-F8-1 construct containing both the light and heavy chains of factor VIII was constructed as follows. A PCR fragment, Z8, containing cloning sites, 5' - splicing donor site of a synthetic intron based on EF1 α and immunoglobulin G (IgG) intron sequences, Kozak sequence and the first 16 nucleotides of the human blood coagulation factor VIII (FVIII) coding sequence was generated using oligonucleotides Z8S and Z8A. The sequences of the nucleic acids is shown below:

Oligonucleotide Z8S:

5' cccaagcttgcggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtgttcc 3'
(SEQ ID NO:1)

Oligonucleotide Z8A:

5' ccgctcgagcagagctctatttgcattggtggaatcgatgccgcggaaccacacacggc 3'
(SEQ ID NO:2)

PCR fragment Z8:

5' cccaagcttgcggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtgttcccgccgcatcgattccaccatgcaaataagctctgctcgagcgg 3' (SEQ ID NO:3)

Nucleic acid Z8 was inserted into pZERO-2 (Invitrogen) between *HindIII* and *XhoI* sites to create pZ8. A PCR fragment, INT3, containing the branching point, the polypyrimidine tract, and the 3' splicing acceptor site of the synthetic intron was generated using oligonucleotides INT3S and INT3A whose sequence is shown below.

Oligonucleotide INT3S:

5' ttcccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactga 3'
(SEQ ID NO:4)

Oligonucleotide INT3A:

5' gaatcgataacctgtggagaaaaagaaaaagtggatgtcagtgtaattcaaggc 3'
(SEQ ID NO:5)

PCR Fragment INT3:

5' ttcccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactgacact
gacatccactttttcttttctccacaggtatcgattc 3' (SEQ ID NO:6)

INT3 was inserted between the *SacII* and *ClaI* sites of pZ8 to create pZ8.I. Therefore, Z8.I contains the entire synthetic intron between *AvrII* and *ClaI* sites. A hFVIII cDNA fragment having *SacI* and *XhoI* restriction sites was inserted between the *SacI* and *XhoI* sites of pZ8.I to create pZ8.I.dB. Therefore, pZ8.I.dB contains a synthetic intron and the entire coding sequence of hFVIII.

A PCR fragment, EG3, containing three HNF-3 binding sites and -54 to +8 of mouse albumin gene was generated using oligonucleotides EG3S and EG3A with modifications to eliminate linker sequences. The sequences of EG3S and EG3A are as follows:

Oligonucleotide EG3S:

5' aggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaatacca
tctacagttattgggttaa 3' (SEQ ID NO:7)

Oligonucleotide EG3A:

5' ggaaaggtgatctgtgtgcagaaagactcgctctaatacttctttaaccaataactg 3'

5 (SEQ ID NO:8)

PCR Fragment EG3:

5' agggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaataccatccaggggaatgtttgttctta
aataccatctacagttattggtaaagaagtattagagcgagtccttctgcacacagatcacctttcc 3' (SEQ

10 ID NO:9)

EG3 was then phosphorylated using T4 polynucleotide kinase and inserted into the *Sma*I site of pZ8.I.dB to create pZ8.I.dB.egg. A DNA fragment, SPA, containing an efficient synthetic polyA signal based on rabbit β -globin sequences (Genes and Develop., 3:1019) was generated by hybridizing two oligonucleotides SPA.S and SPA.A.

15

Oligonucleotide SPA.S:

5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgc 3'

(SEQ ID NO:10)

20

Oligonucleotide SPA.A:

5' tcgagcggccgcacacaaaaaccaacacacagatctctagagctctgatcttttattc 3'

(SEQ ID NO:11)

25 PCR Fragment SPA:

5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgcgcga 3'

(SEQ ID NO:12)

SPA was inserted into the *Xho*I site of pZero-2 to create pZero-2.SPA. SPA was excised from a pZero-2.SPA clone and inserted into the *Xho*I site of pZ8.I.dB.egg to create

30

pZ8.I.dB.egg.A. pAAV-CMV-FIX9 was digested with *Cla*I, blunted with T4 polymerase and religated to create pAAV(*Cla*⁻)-CMV-FIX9.

The entire expression cassette containing HNF-3.albumin promoter-synthetic intron-hFVIII-synthetic poly A signal was excised from pZ8.I.dB.egg.A using *Not*I and ligated to the plasmid backbone and AAV ITRs from pAAV (*Cla*⁻)-CMV-FIX9 to create pAAV-F8-1. The nucleotide sequence of the vector from ITR to ITR (*i.e.*, excluding plasmid backbone) is shown in SEQ ID NO 13.

EXAMPLE 3

Virion Production

AAV vectors were produced from these plasmids using the Ad free system as previously described in U.S. Patent No. 5,858,351; U.S. Patent No. 5,846,528; U.S. Patent No. 5,622,856; and Matsushita *et al.*, Gene Ther 5:938 (1998) all of which are hereby incorporated by reference. Briefly, 293 cells (ATCC, catalog number CRL-1573) were seeded in 10 cm dishes at a density of 3×10^6 cells per dish in 10 ml medium and incubated at 37°C with CO₂ and humidity. After an overnight incubation, cells were approximately seventy to eighty percent confluent.

The cells were then transfected with DNA by the calcium phosphate method, as is well known in the art. Briefly, 7 µg of AAV vector containing the Factor VIII coding region, 7 µg of pladen05 which supplies the accessory functions, and 7 µg of 1909 AAV helper were added to a 3 ml sterile, polystyrene snap cap tube using sterile pipette tips. Then, 1.0 ml of 300 mM CaCl₂ (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na₂PO₄, 12 mM dextrose, pH 7.05, JRH grade) was added with a 2 ml pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the cells, one drop at a time, evenly throughout the 10 cm dish. The cells were then incubated at 37°C with CO₂ and humidity for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were provided with fresh medium and incubated for 72 hours.

After 72 hours, the cells were harvested, pelleted, and resuspended in 1 ml TBS/1% BSA. Freeze/thaw extracts were prepared by repeatedly (three times) freezing the cell suspension on dry ice and thawing at 37°C. Viral preps were stored at -80°C and titered by dot blot assay prior to the first round of infection.

5

EXAMPLE 4

In Vitro Cell Transduction

Cells from the stable human cell line, 293 (ATCC No. CRL1573) were seeded in six-well plates (*i.e.*, plates having six wells for cell growth) at a density of 5×10^5 cells/well. When the monolayers reached 80-90% confluence, they were infected with rAAV virions AAV-e δ D-FVIII-HC, AAV-e δ D-FVIII-LC, an equal ratio of AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC, or AAV-e δ D-FIX at MOIs of 3×10^3 and 3×10^4 . Eighteen hours post infection, the media were replaced with DMEM/10% heat inactivated FBS. The media were collected later for analysis by ELISA (as described below) for FVIII light chain antigen levels, and by the ChromZ FVIII as coagulation activity (FVIII-c) assay (Helena Labs) for biological activity, using the manufacturer's instructions and as described in Example 6.

10

15

EXAMPLE 5

Single Chain Factor VIII Infectivity Assay

20

25

30

In this Example, the infectivity of single chain Factor VIII was investigated. To determine the infectivity of rAAV-hF8-1, HepG2, 293, and H2.35 cells were infected with rAAV-hF8-1 and a control vector rAAV-hF8L at an MOI of 1×10^4 viral particles per cell. Recombinant AAV DNA in infected cells was isolated by Hirt extraction and run on an alkaline agarose gel. Southern blot analysis using an human F8 probe showed that similar amounts of rAAV-hF8-1 and rAAV-hF8L were isolated from uncoated virus in the infected cells. An infectious center assay (ICA) known in the art (*See e.g.*, Snyder, *Current Protocols in Genetics*, Chapter 12, John Wiley & Sons [1997]) was used to further characterize the infectivity of rAAV-hF8-1. In this assay, the infectious particle to total particle ratio of rAAV-F8-1 and that of a control rAAV vector with the genome size of

4645 nucleotides was determined. The results indicated that rAAV-hF8-1 had an infectious particle to total particle ratio that was comparable to the control vector at approximately 1:1000. Taken together, these results indicate rAAV-hF8-1 has similar infectivity as rAAV vectors with the genome size of wild-type AAV.

5

EXAMPLE 6

Factor VIII Protein Expression Assay

An ELISA specific for the light chain of FVIII was used to determine FVIII light chain antigen levels in the 293 cells, as well as the injected animals (described below).
10 NUNC Maxisorb 96 well plates were coated with 50 μ l of a 1:500 dilution of the light chain specific antibody, N77110 (Biodesign International) in a coating buffer overnight at 4°C. The plate was washed three times with wash buffer (PBS, 0.05% Tween 20) and blocked with 200 μ l blocking buffer (PBS, 10% horse serum, 0.05% Tween 20) at room temperature for 1 hour. The plate was washed three times and standards and samples were
15 applied. Bioclone recombinant human FVIII (Baxter Hyland) was used as the standard, and was diluted in blocking buffer to concentrations ranging from 320 ng/ml to 10 ng/ml.

For analysis of transduced culture supernatants, the standards contained 50% media, and for analysis of mouse plasma, the standards were diluted into 10%, in normal pooled mouse plasma (Sigma). A standard assay reference plasma (SARP; Helena Labs)
20 was also included in the assay. Following the loading of the standards and samples (95 μ l/well), the plate was incubated at room temperature for 2 hours, and washed five times with wash buffer (200 μ l/well). A 1:200 dilution of a horseradish peroxidase-conjugated light chain specific antibody, ESH8-HRP, (American Diagnostica) was added (100 μ l/well), and the plate was incubated for 1 hour at room temperature. The plates were then
25 washed four times with wash buffer, and the antigen was detected using an ABTS peroxidase substrate kit (BioRad) according to the manufacturer's instructions. The results are shown in Table 1 of Example 7, below.

EXAMPLE 7

Factor VIII Biological Activity Assay

30

The ChromZ FVIII coagulation activity (FVIII-c) assay (Helena Labs, Beaumont, TX) was used to detect biologically active FVIII in the 293 cells infected as described in Example 4. Bioclone recombinant human FVIII (Baxter Hyland) was used as a standard to analyze transfected culture supernatants. The standards were diluted in plasma dilution buffer (supplied in kit) in the range of 10 ng/ml to 0.313 ng/ml, and were made 2.5% in media. Because this assay can detect both human and murine FVIII activity, it was modified to deplete biologically active human Factor VIII in the mouse plasma. Mouse plasma was pre-incubated with an antibody specific for human FVIII prior to performing the assay. The difference in FVIII activity between the untreated plasma sample and the antibody treated sample represent the amount of biologically active human FVIII in the plasma. The standard used in the assay was normal pooled human plasma (FACT; obtained from George King Biomedical). Serial dilutions of FACT were made in FVIII deficient plasma from undiluted (200 ng/ml) to 6.25 ng/ml. The standards (10 μ l) were incubated at 37°C for 15 min., with or without the addition of 2 μ l antibody N77110. Similarly, mouse plasma samples were diluted in FVIII deficient plasma and 10 μ l of these diluted samples were incubated with or without 2 μ l of N77110 at 37°C for 15 min., and immediately placed on ice. Thus, all incubations with antibody were done in a background of 100% plasma. The antibody adsorbed and non-adsorbed FACT standards, as well as the mouse plasma samples were diluted 1:20 in plasma detection buffer provided in the ChromZ kit. Thus, the final concentration of the FACT standards used in the assay ranged from 10 ng/ml to 0.313 ng/ml.

Twenty five microliters of these dilutions were added to a chilled 96 well plate. With the plate on ice, 25 μ l of FIXa reagent and 50 μ l of FX were added, and the plate was incubated at 37°C for 15 min. Substrate (50 μ l) was added and the plate was incubated for an additional 3 min at 37°C. The reaction was stopped with the addition of 25 μ l 50% acetic acid and the optical density at 405 nm was measured.

As shown below in Table 1, infection of 293 cells with AAV-e δ D-FVIII-HC resulted in no antigen production, as well as no biologically-active protein. Cells infected with AAV-e δ D-FVIII-LC produced FVIII light chain, but no biologically active protein. However, cells transduced with both vectors produced FVIII light chain and biologically

active FVIII in a dose-dependent manner. Transduction of cells with the negative control vector, AAV-e δ D-FIX, resulted in no antigen nor any biologically active FVIII. It was assumed that equal amounts of heavy and light chains were produced in transduced cells. The activity units were converted to nanograms using the definition of one unit being equal to the amount of FVIII in 1 ml of normal pooled human plasma, or 200 ng.

TABLE 1. *In Vitro* Production of Biologically Active Human Factor VIII From Two rAAV Vectors

Vector	MOI	ELISA (ng/ml)	ChromZ	
			(mU/ml)	(ng/ml)
AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC	3×10^3	24	35	7.1
AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC	3×10^4	121	440	87.9
AAV-e δ D-FVIII-HC	3×10^3	0	0	0
AAV-e δ D-FVIII-HC	3×10^4	0	0	0
AAV-e δ D-FVIII-LC	3×10^3	20.5	0	0
AAV-e δ D-FVIII-LC	3×10^4	96.9	0	0
AAV-hFIX	3×10^3	0	0	0
AAV-hFIX	3×10^4	0	0	0
No Vector		0	0	0

Immunofluorescent Staining of FVIII Heavy and Light Chains

In these experiments, 293 cells transduced as described above were analyzed using immunofluorescent staining. 293 cells were plated on rat tail collagen-coated two-well culture slides at a density of 4×10^5 cells per well. Forty-eight hours later, the cells were

transduced at an MOI of 3×10^4 particles per cell of rAAV-hFVIII-HC and rAAV-hFVIII-LC. Forty-eight hours post-transduction, the cells were fixed *in situ* with acetone, blocked with 2% BSA, and stained with a fluorescently labelled anti-hFVIII light chain antibody and a fluorescently labelled anti-hFVIII heavy chain antibody. The anti-hFVIII light chain antibody used was ESH-4 monoclonal antibody (American Diagnostica), fluorescently labelled with alexa-488 (Molecular Probes), according to the manufacturer's instructions. The anti-hFVIII heavy chain antibody used was MAS530P monoclonal antibody (Accurate Chemical) fluorescently labelled with alexa-594 (Molecular Probes), according to the manufacturer's instructions. The cells were counter-stained with DAPI. The images were collected using a Zeiss Axioskop fluorescence microscope equipped with separate filters for DAPI, FITC, and rhodamine signals and a CCD camera. Image analysis was performed using Quips imaging software (Vysis).

As indicated above, infection of cells with either rAAV-e δ D-FVIII-HC or AAV-e δ D-FVIII-LC, followed by staining with antibodies to both chains resulted in production of the individual chains of human FVIII. Immunofluorescent staining of cells co-infected with both vectors demonstrated that although some cells express only the heavy or light chain of hFVIII, many co-expressed both chains of human FVIII.

EXAMPLE 9

In Vitro Expression of Factor VIII Using Single Construct

Table 2 shows that two single vector constructs containing the heavy and the light chain of Factor VIII driven by different promoters express biologically active Factor VIII. The constructs pAAV-hF8-1 (SEQ ID NO:13), and pVm4.1cF8 Δ B (SEQ ID NO:14) were transfected into 293 cells. Following transfection, the cells were allowed to express factor VIII for 48-72 hours. Factor VIII in the culture media was assayed by the ChromZ FVIII coagulation activity (FVIII-c) assay, as per the manufacturer's instructions.

TABLE 2***In Vitro* Production of Biologically Active FVIII**

Construct(s)	ELISA (ng/ml)	ChromZ (ng/ml)
Control	-	0
pAAV-hF8-1	-	4.9
pVm4.1cF8ΔB	-	46

EXAMPLE 10**Factor VIII Expression Using Tissue Specific Promoters**

5 In these experiments, different promoters and enhancer elements were used to drive expression of a Factor VII coding sequence. Expression of Factor VIII was compared in 293 cells and HepG2 cells using different promoters. The pAAVeF8ΔB contains an EF-1α promoter with a hGH intron, Factor VIII with a B-domain deletion (F8ΔB) and a polyA. As described previously, pAAV-hF8-1 uses the HNF-3 albumin promoter with a minimal
10 intron followed by F8ΔB and a minimal polyA. The construct pAAV-c8 uses the CMV enhancer-promoter and the F8ΔB. pAAV8b1 contains the HNF-3 albumin promoter followed by the CMV/B-globin intron with the F8ΔB and a minimal poly A site. Table 3 describes Factor VIII expression using the albumin promoter relative to the control plasmid pV4.1eF8ΔB in HepG2 and 293 cells. These data show increased expression of
15 Factor VIII in HepG2 liver cells with the albumin promoter as compared to Factor VIII expression in 293 cells.

TABLE 3

Relative Tissue Specificity of Promoters

Plasmid Construct	HepG2 Cells	293 Cells
pAAV-hF8-1	6.2	0.6
pAAV8bl	6.7	1.0
pAAVc8	30.0	41.0
pV4.1eF8 Δ B	100	100

Next, several promoters derived from the transthyretin (TTR) gene promoter were transfected into HepG2 cells. TTR is an abundant serum protein and the gene enhancer-promoter contains well known liver-specific transcription factor binding sites (Samadani *et al.*, Gene Expression 6:23 [1996]; Yan *et al.*, EMBO 9:869 [1990]; Costa and Grayson, Nuc. Acids Res., 19:4139 [1991]; Costa *et al.*, Mol. Cell. Bio., 6:4697 [1986]). The constructs were made by replacing the HNF-3 albumin promoter in pAAV-hF8-1 with various lengths of the TTR promoter-enhancer. The TTR enhancer-promoter was modified by replacing the weak affinity binding sites with the strong affinity binding sites to create pAAV-hF8-2. The pAAV-hF8-TTR-E-L-P202 construct contains the full TTR promoter with a linker between the enhancer and the promoter. The remaining constructs are 5' deletions: pAAV-hF8-TTR-E-P202 has the promoter and enhancer with no linker; pAAV-hF8-TTR-E-P197 has a 5 base pair deletion from the promoter; pAAV-hF8-TTR-E-P151 has a 50 base pair deletion; pAAV-hF8-TTR-P202 lacks the TTR enhancer and pAAV-hF8-TTR(X) has a 65 base pair deletion in the enhancer. The control plasmid, pAAV-hF8-1, expressed approximately 4.6 mU/ml. Table 4 shows the fold-increase in Factor VIII activity using the TTR promoter series relative to the control plasmid.

TABLE 4

Factor VIII Expression Using TTR-Derived Promoters

Plasmid Construct	Relative Factor VIII Activity
pAAV-hF8-STTR	3.16
pAAV-hF8-TTR-E-L-P202	8.86
pAAV-hF8-TTR-E-P202	6.1
pAAV-hF8-TTR-EP197	7.3
pAAV-hF8-TTR-E-P151	13.3
pAAV-hF8-TTR-P202	2.3

EXAMPLE 11

In Vivo Expression of Factor VIII

5 In order to test the feasibility of the AAV vector approach of the present invention *in vivo*, three groups of five C57BL/6 mice were injected via the portal vein with either 3×10^{11} particles of AAV-e δ D-FVIII-HC, 3×10^{11} particles of AAV-e δ D-FVIII-LC, or 3×10^{11} particles of both AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC. In addition, a
10 group of four animals was injected with 3×10^{11} particles of AAV-e δ D-FIX. It has been shown that this strain of mice does not elicit an immune response to human FVIII when the gene is delivered to the liver via an adenoviral vector (Connelly *et al.*, Blood 87:4671-4677 [1996]). As indicated by the results shown below, the data obtained during these experiments demonstrate the feasibility of producing biologically active FVIII using two
15 AAV vectors to independently deliver the heavy and light chains of FVIII.

Blood samples were collected in sodium citrate via the retro-orbital plexus at biweekly intervals for the first 2 months and at monthly intervals thereafter for 6 months and at 11 months. Very high levels of FVIII light chain were expressed in animals injected with AAV-e δ D-FVIII-LC alone or both vectors as shown in Figure 8.

20 In order to assess the amount of biologically active human FVIII produced in the animals, a modified ChromZ assay was used. Since this assay detects both human and

murine FVIII, the amount of FVIII present in the plasma before and after adsorption to an antibody specific to human FVIII was determined. The amount of FVIII remaining in the plasma after adsorption represented the amount of active murine FVIII and the difference represented the amount of active human FVIII. Control experiments demonstrated that the antibody could remove 80-90% of the human FVIII from a mouse plasma sample when the sample was spiked with up to 32 ng of human FVIII. The modified ChromZ assay indicated that only those animals injected with both vectors produced biologically active FVIII, as indicated in Table 5. The results shown in Table 5 are those from plasma collected 8 weeks post-injection, although similar results were obtained at 10 weeks and 5 months post-injection. One of the five animals co-injected with both the heavy and light chain vector did not express VIII, presumably due to an inefficient injection, and was omitted from the analysis. Animals injected with both vectors produced over 2 µg/ml hFVIII light chain as measured by ELISA. The ChromZ assay indicated that a total of 600-900 ng/ml of active hFVIII was detected in the plasma. The contribution from murine Factor VIII was approximately 400-500 ng/ml, indicating that about 230-430 ng/ml of active human Factor VIII was present in the plasma. Although only a fraction of the total protein was found to be active, the animals produced physiological levels of the active protein (*i.e.*, 200 ng/ml). The animals were found to have maintained these physiological levels of active protein for more than 11 months, without waning. Similar analyses performed on animals injected with the light chain vector alone, the heavy chain vector alone, or the hFIX vector demonstrated no biologically active human FVIII in the plasma of these animals.

TABLE 5

Biological Activity of Human Factor VIII *In Vivo*

Construct(s) Used	ELISA (ng/ml)	Total FVIII (-Ab) (Units)	Murine FVIII (+Ab) (Units)	Human FVIII (ng/ml)
AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC*	2288	3.9	2.2	342
AAV-e δ D-FVIII-LC *	3329	1.4	1.6	0
AAV-e δ D-FVIII-HC	0	1.6	1.6	0
AAV-e δ D-FIX	0	1.4	2.0	0

*Average of three animals.

5

EXAMPLE 12

Gene Transfer and Vector Expression in Tissues

In these experiments, evidence of gene transfer to liver was obtained by Southern Blot analysis following isolation of DNA from one animal of each experimental group sacrificed 8 weeks post-injection (*i.e.*, as described in Example 11). In addition, DNA was obtained from other tissues in order to determine the degree of vector expression in organs other than the liver.

10

15

20

Twenty micrograms of DNA was digested with *Bgl*II, separated using a 1% agarose gel, and hybridized with a ³²P-labelled 1126 bp *Alw*NI fragment encoding the A1 and A2 domains of hFVIII (heavy chain probe), or a ³²P-labelled 1456 bp *Nde*I-*Eco*RI fragment encoding the A3, C1 and C2 domains of hFVIII (light chain probe). Copy number controls were generated by spiking *Bgl*II-digested naive mouse liver DNA with *Bgl*II-digested heavy or light chain plasmid DNA (pVm4.1e δ D-hFVIII-HC and pVm4.1e δ D-hFVIII-LC, respectively), at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome. The hybridized membranes were analyzed using a Storm 860 phosphorimager (Molecular Dynamics), and quantitation of vector copy number was evaluated using ImageQuaNT software (Molecular Dynamics). Autoradiography of the hybridized membranes was also performed. Total

RNA was isolated from liver tissue using the RNA Stat extraction kit (Tel-Test). As describe briefly below, Northern blot analysis was also performed on 10 µg RNA using methods known in the art, in conjunction with the ³²P-labelled probes specific to the heavy and light chains of hFVIII described above and autoradiography was performed on the hybridized membranes.

Following digestion of liver DNA with *Bgl*II and hybridization with an hFVIII light chain probe described below, using methods known in the art, a band at the predicted size of 3015 bp was detected in animals injected with rAAV-hFVIII-LC, or both the heavy and light chain vectors. This band was not observed in the DNA of animals injected with the heavy chain vector alone or the hFIX vector, as shown in Figure 9A (rAAV-hFVIII-LC, lane 1; rAAV-hFIX, lane 2; rAAV-hFVIII-HC, lane 3; both rAAV-hFVIII-LC and rAAV-hFVIII-HC, lane 4; copy number controls were generated by spiking *Bgl*II digested naive mouse liver DNA with the corresponding plasmids at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome, lanes 5-9). Phosphoimage analysis revealed that the light chain vector was present at approximately 2.4 and 1.5 copies per diploid genome in animals injected with the light chain vector alone or both vectors, respectively. When *Bgl*II-digested DNA was hybridized with an hFVIII heavy chain probe, the expected band of 2318 bp was observed in animals injected with the heavy chain vector alone or both vectors, but was not detected in animals injected with the light chain vector alone or the hFIX vector, as shown in Figure 9B. The copy number in animals injected with the heavy chain vector alone and both vectors was 1.1 and 1.7 vector copies per diploid genome, respectively.

The results of hybridization of DNA extracted from the spleen, kidney and heart tissue with either an hFVIII light chain probe or a heavy chain probe indicated that these tissues contained less than 1 copy of vector sequences per 10 diploid genomes, demonstrating that the vector distributes primarily to the liver following intra-portal injection, as shown in Figures 10A and 10B.

Human FVIII gene expression in the liver of the mice was also assessed by Northern blot analysis on RNA isolated from animals sacrificed 8 weeks post-injection (as described above). hFVIII light chain transcripts of the predicted size (2.7 kb) were

observed in animals injected with the light chain vector alone or both vectors, as shown in Figure 11A. Similarly, the expected hFVIII heavy chain transcripts (2.7 kb) were detected in animals that were injected with the heavy chain vector alone or both vectors, as shown in Figure 11B. Since the heavy and light chain DNA sequences were shown by Southern blot analysis to be present at approximately the same copy number (1.7 and 1.5 copies per diploid genome, respectively), in an animal injected with both vectors, these results demonstrate that both the heavy and light chains of hFVIII are expressed in the liver in approximately equivalent amounts.

EXAMPLE 13

In vivo Expression of Factor VIII Using Single Vectors

Several groups comprising four C57BL/6 mice were injected via the portal vein with 3×10^{11} particles of AAV-hF8-STTR, 3×10^{11} particles of AAV-hF8-hAAT-137, or 3×10^{11} particles of AAV-hF8-HNF3-alb-original. As discussed above, AAV-hF8-STTR contains modified sequences from the transthyretin gene promoter. AAV-hF8-hAAT-137 contains 137 base pairs of the human₁-antitrypsin promoter. See De Simone et al., EMBO J., 6:2759-2766 (1987). AAV-hF8-HNF3-alb-original, like AAV-F8-1, contains three HNF3 binding sites and 54 base pairs of the albumin promoter.

Expression of Factor VIII was measured at 4 weeks by human Factor VIII-specific ELISA (Affinity Biologicals). Even at this early time-point, several of these mice expressed between 2 and 20 ng/ml of human Factor VIII. Table 6 shows expression levels at 4 weeks post-infection for selected animals.

TABLE 6. Factor VIII Expression at 4 Weeks

rAAV	Factor VIII Levels (ng/ml)
rAAV-hF8-STTR (mouse #1)	4.3
rAAV-hF8-STTR (mouse #2)	2.4
rAAV-hF8-hAAT-137 (mouse #1)	10.5

rAAV-hF8-hAAT-137 (mouse #2)	20.4
rAAV-hF8-HNF3-alb-original (mouse #1)	2.2

rAAV-hF8-hAAT-137 produced expression levels of as much as 10 to 20 ng/ml, which represent 20% to 40% of normal human levels of Factor VIII. These data therefore show the *in vivo* expression of therapeutically effective amounts of human Factor VIII using

SEQUENCE LISTING

5 <110> Couto, Linda B.
 Colosi, Peter C.
 <120> Adeno-Associated Vectors for Expression of Factor VIII
 by Target Cells
 <130> Avigen-04082
 10 <140> xx/xxx,xxx
 <141> 2000-12-18
 <150> 09/470,618
 15 <151> 1999-12-22
 <150> 09/364,862
 <151> 1999-07-30
 <150> 60/125,974
 20 <151> 1999-03-24
 <150> 60/104,994
 <151> 1998-10-20
 25 <160> 15
 <170> PatentIn Ver. 2.0
 <210> 1
 30 <211> 59
 <212> DNA
 <213> Artificial Sequence
 <220>
 35 <223> Description of Artificial Sequence: Synthetic
 <400> 1
 cccaagcttg cggccgcccg ggtgccgccc ctaggcaggt aagtgccgtg tgtggttcc 59
 40 <210> 2
 <211> 59
 <212> DNA
 <213> Artificial Sequence
 45 <220>
 <223> Description of Artificial Sequence: Synthetic

<400> 2
 ccgctcgagc agagctctat ttgcatggtg gaatcgatgc cgcgggaacc acacacggc 59

5 <210> 3
 <211> 103
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence: Synthetic

15 <400> 3
 cccaagcttg cggccgccc ggtgccgcc ctaggcaggt aagtgccgtg tgtggttccc 60
 gcggcatcga ttccaccatg caaatagagc tctgctcgag cgg 103

20 <210> 4
 <211> 57
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> Description of Artificial Sequence: Synthetic

30 <400> 4
 ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactga 57

35 <210> 5
 <211> 57
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence: Synthetic

45 <400> 5
 gaatcgatac ctgtggagaa aaagaaaaag tggatgtcag tgtcagtaat tcaaggc 57

40 <210> 6
 <211> 99
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> Description of Artificial Sequence: Synthetic

45 <400> 6
 ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactgacac 60
 tgacatccac tttttctttt tctccacagg tatcgattc 99

<210> 7
 <211> 100
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 7
 10 agggaatggtt tggtcttaaa taccatccag ggaatgtttg ttcttaaata ccattccaggg 60
 aatgtttggtt cttaaatacc atctacagtt attggttaaa 100
 <210> 8
 <211> 59
 15 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 20 <400> 8
 ggaaagggtga tctgtgtgca gaaagactcg ctctaataata cttctttaac caataactg 59
 <210> 9
 25 <211> 144
 <212> DNA
 <213> Artificial Sequence
 <220>
 30 <223> Description of Artificial Sequence: Synthetic
 <400> 9
 agggaatggtt tggtcttaaa taccatccag ggaatgtttg ttcttaaata ccattccaggg 60
 aatgtttggtt cttaaatacc atctacagtt attggttaaa gaagtattatt agagcgagtc 120
 35 tttctgcaca cagatcacct ttcc 144
 <210> 10
 <211> 59
 <212> DNA
 40 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 10
 45 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgc 59
 <210> 11

<211> 59
 <212> DNA
 <213> Artificial Sequence

5

<220>
 <223> Description of Artificial Sequence: Synthetic

<400> 11
 tcgagcggcc gcacacaaaa aaccaacaca cagatctcta gagctctgat cttttattc 59

10

<210> 12
 <211> 63
 <212> DNA
 <213> Artificial Sequence

15

<220>
 <223> Description of Artificial Sequence: Synthetic

<400> 12
 20 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgct 60
 cga 63

<210> 13
 <211> 11933
 25 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

30

<400> 13
 cagctgcgcg ctgcctcgct cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc 60
 tttggctgcc cggcctcagt gagcgagcga gcgcgcagag agggagtggc caactccatc 120
 actaggggtt cctgcggccg cccagggaat gtttgttctt aaataccatc cagggaatgt 180
 35 ttgttcttaa ataccatcca gggaatgttt gttcttaaata accatctaca gttattgggt 240
 aaagaagtat attagagcga gtctttctgc acacagatca cctttccggg tgcgcgccct 300
 aggcaggtaa gtgccgtgtg tggttccgcg gggcctggcc tctttacggg ttatggccct 360
 tgcgtgcctt gaattactga cactgacatc cactttttct ttttctccac aggtatcgat 420
 tccaccatgc aaatagagct ctccacctgc ttctttctgt gccttttgcg attctgcttt 480
 40 agtgccacca gaagatacta cctgggtgca gtggaactgt catgggacta tatgcaaagt 540
 gatctcgggtg agctgcctgt ggacgcaaga tttcctccta gagtgccaaa atcttttcca 600
 ttcaacacct cagtcgtgta caaaaagact ctgttttag aattcacgga tcaccttttc 660
 aacatcgcta agccaaggcc accctggatg ggtctgctag gtccctaccat ccaggctgag 720
 gtttatgata cagtggatcat tacacttaag aacatggctt cccatcctgt cagtcttcat 780
 45 gctgttggtg tctcctactg gaaagcttct gagggagctg aatatgatga tcagaccagt 840
 caaagggaga aagaagatga taaagtcttc cctgggtgaa gccatacata tgtctggcag 900
 gtccctgaaag agaatggtcc aatggcctct gaccactgt gccttaccta ctcatatctt 960
 tctcatgtgg acctggtaaa agacttgaat tcaggcctca ttggagccct actagtatgt 1020

	agagaaggga	gtctggccaa	ggaaaagaca	cagaccttgc	acaaatttat	actacttttt	1080
	gctgtatttg	atgaaggga	aagttggcac	tcagaaacaa	agaactcctt	gatgcaggat	1140
	agggatgctg	catctgctcg	ggcctggcct	aaaatgcaca	cagtcaatgg	ttatgtaaac	1200
	aggtctctgc	caggtctgat	tggatgccac	aggaaatcag	tctattggca	tgtgattgga	1260
5	atgggcacca	ctcctgaagt	gcactcaata	ttcctcgaag	gtcacacatt	tcttgtgagg	1320
	aaccatcgcc	aggcgtcctt	ggaaatctcg	ccaataactt	tccttactgc	tcaaacactc	1380
	ttgatggacc	ttggacagtt	tctactgttt	tgtcatatct	cttcccacca	acatgatggc	1440
	atggaagctt	atgtcaaagt	agacagctgt	ccagagggaac	cccaactacg	aatgaaaaat	1500
	aatgaagaag	cggaagacta	tgatgatgat	cttactgatt	ctgaaatgga	tgtggtcagg	1560
10	tttgatgatg	acaactctcc	ttcctttatc	caaattcgct	cagttgccaa	gaagcatcct	1620
	aaaacttggg	tacattacat	tgctgctgaa	gaggaggact	gggactatgc	tcccttagtc	1680
	ctcgcccccg	atgacagaag	ttataaaagt	caatatttga	acaatggccc	tcagcggatt	1740
	ggtaggaagt	acaaaaaagt	ccgattttatg	gcatacacag	atgaaacctt	taagactcgt	1800
	gaagctattc	agcatgaatc	aggaatcttg	ggacctttac	tttatgggga	agttggagac	1860
15	acactgttga	ttatatttaa	gaatcaagca	agcagaccat	ataacatcta	ccctcacgga	1920
	atcactgatg	tccgtccttt	gtattcaagg	agattaccaa	aagggtgtaa	acatttgaag	1980
	gattttccaa	ttctgccagg	agaaatattc	aaatataaat	ggacagtgc	tgtagaagat	2040
	gggccaacta	aatcagatcc	tcggtgcctg	acccgctatt	actctagttt	cgtaaatatg	2100
	gagagagatc	tagcttcagg	actcattggc	cctctcctca	tctgctacaa	agaatctgta	2160
20	gatcaaaag	gaaaccagat	aatgtcagac	aagagggaatg	tcacctctgt	ttctgtattt	2220
	gatgagaacc	gaagctggta	cctcacagag	aatatacaac	gctttctccc	caatccagct	2280
	ggagtgcagc	ttgaggatcc	agagttccaa	gcctccaaca	tcatgcacag	catcaatggc	2340
	tatgtttttg	atagtttgca	gttgtcagtt	tgtttgcatg	aggtggcata	ctggtacatt	2400
	ctaagcattg	gagcacagac	tgacttcctt	tctgtcttct	tctctggata	taccttcaaa	2460
25	cacaaaatgg	tctatgaaga	cacactcacc	ctattcccat	tctcaggaga	aactgtcttc	2520
	atgtcgaatg	aaaaccagg	tctatggatt	ctgggggtgc	acaactcaga	ctttcggaac	2580
	agaggcatga	ccgccttact	gaaggtttct	agttgtgaca	agaacactgg	tgattattac	2640
	gaggacagtt	atgaagatat	ttcagcatac	ttgctgagta	aaaacaatgc	cattgaacca	2700
	agaagcttcg	aaataactcg	tactactctt	cagtcagatc	aagaggaaat	tgactatgat	2760
30	gataccatat	cagttgaaat	gaagaaggaa	gattttgaca	tttatgatga	ggatgaaaat	2820
	cagagccccc	gcagctttca	aaagaaaaca	cgacactatt	ttattgctgc	agtggagagg	2880
	ctctgggatt	atgggatgag	tagctcccca	catgtttctaa	gaaacagggc	tcagagtggc	2940
	agtgctccctc	agttcaagaa	agttgttttc	caggaattta	ctgatggctc	ctttactcag	3000
	cccttataacc	gtggagaact	aatgaacat	ttgggactcc	tggggccata	tataagagca	3060
35	gaagttgaag	ataatatcat	ggtaactttc	agaaatcagg	cctctcgtcc	ctattccttc	3120
	tattctagcc	ttatttctta	tgaggaagat	cagaggcaag	gagcagaacc	tagaaaaaac	3180
	tttgtcaagc	ctaataaagc	caaaacttac	ttttggaaag	tgcaacatca	tatggcacc	3240
	actaaagatg	agtttgactg	caaagcctgg	gcttatttct	ctgatgttga	cctggaaaaa	3300
	gatgtgcact	caggcctgat	tggacccctt	ctggtctgcc	acactaacac	actgaaccct	3360
40	gctcatggga	gacaagtgc	agtacaggaa	tttgctctgt	ttttcaccat	ctttgatgag	3420
	acaaaaagct	ggtacttcac	tgaaaaatatg	gaaagaaaact	gcagggctcc	ctgcaatatc	3480
	cagatggaag	atcccacttt	taaagagaat	tatcgcttcc	atgcaatcaa	tggctacata	3540
	atggatacac	tacctggctt	agtaatggct	caggatcaaa	ggattcgatg	gtatctgctc	3600
	agcatgggca	gcaatgaaaa	catccattct	attcatttca	gtggacatgt	gttactgta	3660
45	cgaaaaaaag	aggagtataa	aatggcactg	tacaatctct	atccagggtg	ttttgagaca	3720
	gtggaaatgt	taccatccaa	agctggaatt	tggcgggtgg	aatgccttat	tggcgagcat	3780
	ctacatgctg	ggatgagcac	actttttctg	gtgtacagca	ataagtgtca	gactcccctg	3840
	ggaatggctt	ctggacacat	tagagatttt	cagattacag	cttcaggaca	atatggacag	3900

	tgggccccaa	agctggccag	acttcattat	tccggatcaa	tcaatgcctg	gagcaccaag	3960
	gagccctttt	cttggatcaa	ggtggatctg	ttggcaccac	tgattattca	cggcatcaag	4020
	acccaggggtg	cccgtcagaa	gttctccagc	ctctacatct	ctcagtttat	catcatgtat	4080
	agtcttgatg	ggaagaagtg	gcagacttat	cgaggaaatt	ccactggaac	cttaatgggtc	4140
5	ttctttggca	atgtggattc	atctgggata	aaacacaata	tttttaaccc	tccaattatt	4200
	gctcgatata	tccgtttgca	cccaactcat	tatagcattc	gcagcactct	tcgcatggag	4260
	ttgatgggct	gtgattttaa	tagttgcagc	atgccattgg	gaatggagag	taaagcaata	4320
	tcagatgcac	agattactgc	ttcatcctac	tttaccaata	tgtttgccac	ctgggtctct	4380
	tcaaaagctc	gacttcacct	ccaagggagg	agtaatgcct	ggagacctca	ggtgaataat	4440
10	ccaaaagagt	ggctgcaagt	ggacttccag	aagacaatga	aagtcacagg	agtaactact	4500
	cagggagtaa	aatctctgct	taccagcatg	tatgtgaagg	agttcctcat	ctccagcagt	4560
	caagatggcc	atcagtggac	tctctttttt	cagaatggca	aagtaaaggt	ttttcagggg	4620
	aatcaagact	ccttcacacc	tgtggtgaac	tctctagacc	caccgttact	gactcgctac	4680
	cttcgaattc	acccccagag	ttgggtgcac	cagattgccc	tgaggatgga	ggttctgggc	4740
15	tgcgaggcac	aggacctcta	ctgactcgag	aataaaagat	cagagctcta	gagatctgtg	4800
	tgttggtttt	ttgtgtgcgg	ccgcaggaac	ccctagtgat	ggagtgggcc	actccctctc	4860
	tgcgcgctcg	ctcgctcact	gaggccgggc	gaccaaaggt	cggccgacgc	ccgggctttg	4920
	cccgggcggc	ctcagtgagc	gagcgagcgc	gcagctgcct	gcaggacatg	tgagcaaaag	4980
	gccagcaaaa	ggccaggaac	cgtaaaaagg	ccgcgttgct	ggcgtttttc	cataggctcc	5040
20	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	gaggtggcga	aaccgcacag	5100
	gactataaag	ataccaggcg	tttccccctg	gaagctccct	cgtgcgctct	cctgttccga	5160
	ccctgccgct	taccggatac	ctgtccgcct	ttctcccttc	gggaagcgtg	gcgctttctc	5220
	atagctcacg	ctgtaggtat	ctcagttcgg	tgtaggtcgt	tcgctccaag	ctgggctgtg	5280
	tgcacgaacc	ccccgttcag	cccgaaccgt	gcgccttatc	cggtaactat	cgtcttgagt	5340
25	ccaaccgggt	aagacacgac	ttatcgccac	tggcagcagc	cactggtaac	aggattagca	5400
	gagcgaggta	tgtaggcggt	gctacagagt	tcttgaagtg	gtggcctaac	tacggctaca	5460
	ctagaaggac	agtatttgggt	atctgcgctc	tgtgaagcc	agttaccttc	ggaaaaagag	5520
	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cgggtggttt	tttgtttgca	5580
	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc	ttttctacgg	5640
30	ggtctgacgc	tcagtggaa	gaaaactcac	gttaagggat	tttgggtcatg	agattatcaa	5700
	aaaggatctt	cacctagatc	cttttaaat	aaaaatgaag	ttttaaatca	atctaaagta	5760
	tatatgagta	aacttgggtc	gacagttacc	aatgcttaat	cagtgaggca	cctatctcag	5820
	cgatctgtct	atttcgttca	tccatagttg	cctgactccc	cgtcgtgtag	ataactacga	5880
	tacgggaggg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	5940
35	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	agaagtgggtc	6000
	ctgcaacttt	atccgcctcc	atccagtcta	ttaattgttg	ccgggaagct	agagtaagta	6060
	gttcgccagt	taatagtttg	cgcaacggtg	ttgccattgc	tacaggcatc	gtgggtgtcac	6120
	gctcgtcggt	tggtatggct	tcattcagct	ccggttccca	acgatcaagg	cgagttacat	6180
	gatcccccat	gttgtgcaaa	aaagcgggta	gctccttcgg	tcctccgatc	gttgtcagaa	6240
40	gtaagttggc	cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat	tctcttactg	6300
	tcatgccatc	cgtaagatgc	ttttctgtga	ctgggtgagta	ctcaaccaag	tcattctgag	6360
	aatagtgtat	gcggcgaccg	agttgtctct	gcccggcgtc	aatacgggat	aataccgcgc	6420
	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	cgaaaactct	6480
	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	6540
45	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	aggcaaatg	6600
	ccgcaaaaaa	gggaataaag	gcgacacgga	aatgttgaat	actcatactc	ttcctttttc	6660
	aatattattg	aagcatttat	caggggtatt	gtctcatgag	cggatacata	tttgaatgta	6720
	tttagaaaaa	taaacaaata	ggggttccgc	gcacatttcc	ccgaaaagtg	ccacctgacg	6780

	tctaagaaac	cattattatc	atgacattaa	cctataaaaa	taggcgtatc	acgaggccct	6840
	ttcgtctcgc	gcgtttcggt	gatgacggtg	aaaacctctg	acacatgcag	ctcccggaga	6900
	cggtcacagc	ttgtctgtaa	gcggatgccg	ggagcagaca	agcccgtcag	ggcgcgtcag	6960
	cgggtggttg	cgggtgtcgg	ggctggctta	actatgcggc	atcagagcag	attgtactga	7020
5	gagtgcacca	taaaattgta	aacgttaata	ttttgttaaa	attcgcgtta	aatttttgtt	7080
	aaatcagctc	atTTTTtaac	caataggccg	aaatcggcaa	aatcccttat	aaatcaaaag	7140
	aatagcccga	gatagggttg	agtgttgttc	cagtttgtaa	caagagtcca	ctattaaaga	7200
	acgtggactc	caacgtcaaa	gggcgaaaaa	ccgtctatca	ggcgatggc	ccactacgtg	7260
	aaccatcacc	caaatacaag	tttttggggt	cgaggtgccg	taaagcacta	aatcgggaacc	7320
10	ctaaaggagg	cccccgattt	agagcttgac	ggggaaagcc	ggcgaacgtg	gcgagaaagg	7380
	aagggaagaa	agcgaagga	gcgggcgcta	gggcgtggc	aagtgtagcg	gtcacgctgc	7440
	gcgtaaccac	cacacccgcc	gcgcttaatg	cgccgctaca	gggcgcgtac	tatggttgct	7500
	ttgacgtatg	cgggtgtgaa	taccgcacag	atgcgtaagg	agaaaatacc	gcatacaggcc	7560
	gtaacctgtc	ggatcaccgg	aaaggacccg	taaagtata	atgattatca	tctacatatc	7620
15	acaacgtgcg	tggaggccat	caaaccacgt	caaataatca	attatgacgc	aggtatcgta	7680
	ttaattgatc	tgcatacaact	taacgtaaaa	acaacttcag	acaatacaaa	tcagcgacac	7740
	tgaatacggg	gcaacctcat	gtcaacgaag	aacagaaccc	gcagaacaac	aaccgcgaac	7800
	atccgctttc	ctaaccaaat	gattgaacaa	attaacatcg	ctcttgagca	aaaagggtcc	7860
	gggaatttct	cagcctgggt	cattgaagcc	tgccgtcgga	gactaacgtc	agaaaagaga	7920
20	gcatatacat	caattaaaag	tgatgaagaa	tgaacatccc	gcgttcttcc	ctccgaacag	7980
	gacgatattg	taaattcact	taattacgag	ggcattgcag	taattgagtt	gcagttttac	8040
	cactttcctg	acagtgcacg	actgcgtggt	ggctctgtca	cagactaaat	agtttgaatg	8100
	attagcagtt	atggtgatca	gtcaaccacc	agggaataat	ccttcatatt	attatcgtgc	8160
	ttcaccaacg	ctgcctcaat	tgctctgaat	gcttccagag	acaccttatg	ttctatacat	8220
25	gcaattacaa	catcagggtg	actcatagaa	atggtgctat	taagcatatt	ttttacacga	8280
	atcagatcca	cggagggatc	atcagcagat	tgctctttat	tcattttgtc	gctccatgcg	8340
	cttgctcttc	atctagcggg	taaaatatta	cttcaaactc	ttctgtatga	agatttgagc	8400
	acgttggcct	tacatacatc	tgctcggttg	atctccctcc	agaatgccag	caggaccgca	8460
	ctttgtttacg	caaccaatac	tattaagtga	aaacattcct	aatatttgac	ataaatcatc	8520
30	aacaaaacac	aaggagggtca	gaccagattg	aaacgataaa	aacgataatg	caaactacgc	8580
	gccctcgtat	cacatggaag	gtttttacca	tggctcagggt	tgccattttt	aaagaaatat	8640
	tcgatcaagt	gcgaaaagat	ttagactgtg	aattgtttta	ttctgaacta	aaacgtcaca	8700
	acgtctcaca	ttatatattac	tatctagcca	cagataatat	tcacatcgtg	ttagaaaacg	8760
	ataacaccgt	gttaataaaa	ggacttaaaa	aggttgtaaa	tgtaaatttc	tcaagaaaca	8820
35	cgcattcttat	agaaacgtcc	tatgataggt	tgaaatcaag	agaaatcaca	tttcagcaat	8880
	acagggaaaa	tcttgctaaa	gcaggagttt	tccgatgggt	tacaaatatc	catgaacata	8940
	aaagatatata	ctataccttt	gataattcat	tactattttac	tgagagcatt	cagaacacta	9000
	cacaaatctt	tccacgctaa	atcataacgt	ccggtttctt	ccgtgtcagc	accggggcgt	9060
	tggcataatg	caatacgtgt	acgcgctaaa	ccctgtgtgc	atcgttttta	ttattcccgg	9120
40	acactcccgc	agagaagttc	cccgctcagg	ctgtggacat	agttaatccg	ggaataacaat	9180
	gacgattcat	cgcacctgac	atacattaat	aaatattaac	aatatgaaat	ttcaactcat	9240
	tgtttaggggt	ttgttttaatt	ttctacacat	acgattctgc	gaacttcaaa	aagcatcggg	9300
	aataacacca	tgaaaaaaat	gctactcgct	actgcgctgg	ccctgcttat	tacaggatgt	9360
	gctcaacaga	cgtttactgt	tcaaaaacaaa	ccggcagcag	tagcaccaaa	ggaaaccatc	9420
45	acccatcatt	tcttcgtttc	tggaattggg	cagaagaaaa	ctgtcgatgc	agccaaaatt	9480
	tgtggcggcg	cagaaaaatgt	tgtaaaaaca	gaaacccagc	aaacattcgt	aatgggattg	9540
	ctcggtttta	ttacttttagg	cattttatact	ccgctggaag	cgcgtgtgta	ttgctcaciaa	9600
	taattgcatg	agttgcccac	cgcgatatgg	gcaactctat	ctgcactgct	cattaatata	9660

```

cttctggggt ccttccagtt gtttttgc atgtgatcagc ctctctctga gggtgaaata 9720
atcccgttca gcggtgtctg ccagtcgggg ggaggctgca ttatccacgc cggaggcggg 9780
gggtggcttca cgcactgact gacagactgc tttgatgtgc aaccgacgac gaccagcggc 9840
aacatcatca cgcagagcat ctttttcagc tttagcatca gctaactcct tcgtgtattt 9900
5  tgcattcgagc gcagcaacat cacgctgacg catctgcatg tcagtaattg ccgcgttcgc 9960
cagcttcagt tctctggcat ttttgcgcg ctgggctttg taggtaattg cgttatcacg 10020
gtaatgatta acagcccatg acaggcagac gatgatgcag ataaccagag cggagataat 10080
cgcggtgact ctgctcatac atcaatctct ctgaccgttc cggccgcttc tttgaatttt 10140
gcaatcaggc tgtcagcctt atgctcgaac tgaccataac cagcgcggcg cagtgaagcc 10200
10 cagatattgc tgcaacggtc gattgcctga cggatatcac cacgatcaat cataggtaaa 10260
gcgccacgct ccttaatctg ctgcaatgcc acagcgctct gacttttcgg agagaagtct 10320
ttcaggccaa gctgcttgcg gtaggcatcc caccaacggg aaagaagctg gtagcgtccg 10380
gcgcctgttg atttgagttt tgggttttag gtgacaagtt tgcgaggggtg atcggagtaa 10440
tcagtaaata gctctccgcc tacaatgacg tcataaccat gatttctggt tttctgacgt 10500
15 ccgttatcag ttccctccga ccacgccagc atatcgagga acgccttacg ttgattattg 10560
atttttacca tcttctactc cggctttttt agcagcgaag cgtttgataa gcgaaccaat 10620
cgagtcagta ccgatgtagc cgataaacac gctcgttata taagcgagat tgctacttag 10680
tccggcgaag tccgagaagg cacgaatgaa ccaggcgata atggcgacac tcgttgctgc 10740
gattactggt tttgtaaacg caccgccatt atatctgccg cgaaggtagc ccattgcaaa 10800
20 cgcaaggatt gccccgatgc cttgttcctt tgccgcgaga atggcgccca acaggtcatg 10860
tttttctggc atcttcatgt cttaccccca ataaggggat ttgctctatt taattaggaa 10920
taaggtcgat tactgataga acaaatccag gctactgtgt ttagtaatca gatttggtcg 10980
tgaccgatat gcacgggcaa aacggcgagg ggttggttagc gcgacctcct gccacccgct 11040
ttcacgaagg tcatgtgtaa aaggccgcag cgtaactatt actaatgaat tcaggacaga 11100
25 cagtggctac ggctcagttt ggggttggtg gttgctgggc ggcgatgacg cctgtacgca 11160
tttgggtgat cggttctgct tccggtattc gcttaattca gcacaacgga aagagcactg 11220
gctaaccagg ctgcgcgact cttcacgatt atcgactcaa tgctcttacc tgttgtgcag 11280
atataaaaaa tcccgaacc gttatgcagg ctctaactat tacctgcgaa ctgtttcggg 11340
attgcatttt gcagacctct ctgcctgcga tggttggagt tccagacgat acgtcgaagt 11400
30 gaccaactag gcggaatcgg tagtaagcgc cgctctttt catctcacta ccacaacgag 11460
cgaattaacc catcggttag tcaaatttac ccaattttat tcaataagtc aatatcatgc 11520
cgtaaatatg ttgccatccg tggcaatcat gctgctaacg tgtgaccgca ttcaaaatgt 11580
tgtctgcgat tgactcttct ttgtggcatt gcaccaccag agcgtcatac agcggcttaa 11640
cagtgcgtga ccaggtgggt tgggtaagggt ttgggattag catcgtcaca gcgcgatatg 11700
35 ctgcgcttgc tggcatcctt gaatagccga cgcctttgca tcttccgcac tctttctcga 11760
caactctccc ccacagctct gttttggcaa tatcaaccgc acggcctgta ccatggcaat 11820
ctctgcatct tgccccgggc gtcgcggcac tacggcaata atccgcataa gcgaatgttg 11880
cgagcacttg cagtaccttt gccttagtat ttccttcaag ctgcccctgc agg 11933

```

```

40 <210> 14
    <211> 4999
    <212> DNA
    <213> Artificial Sequence

```

```

45 <220>
    <223> Description of Artificial Sequence: Synthetic

```

```

<400> 14

```

	cgccccctgca	ggcagctgcg	cgctcgctcg	ctcactgagg	ccgcccgggc	aaagcccggg	60
	cgtcgggcca	ccttttggtcg	cccggcctca	gtgagcgagc	gagcgcgag	agagggagtg	120
	gccaactcca	tcactagggg	ttcctgcggc	cgcacgcgtg	gtggcgcggg	gtaaaactggg	180
5	aaagtgatgt	cgtgtactgg	ctccgccttt	ttcccagggg	tgggggagaa	ccgtatataa	240
	gtgcagtagt	cgccgtgaac	gttctttttc	gcaacgggtt	tgccgccccg	cggcaggtaa	300
	gtgccaggga	atgtttgttc	ttaaatacca	tcgtccagg	gaatgtttgt	tcttaaatac	360
	catctactga	cactgacatc	cactttttct	ttttctccac	aggtatcgat	ccaccatgca	420
	aatagagctc	tccacctgct	tctttctgtg	ccttttgcca	ttctgcttta	gtgccaccag	480
10	aagatactac	ctgggtgcag	tggaaactgtc	atgggactat	atgcaaagt	atctcggtga	540
	gctgcctgtg	gaçgcaagat	ttcctcctag	agtgcacaaa	tcttttccat	tcaacacctc	600
	agtcgtgtac	aaaaagactc	tgtttgtaga	attcacggat	caccttttca	acatcgctaa	660
	gccaaggcca	ccctggatgg	gtctgctagg	tcctaccatc	caggctgagg	tttatgatac	720
	agtggtcatt	acacttaaga	acatggcttc	ccatcctgtc	agtcttcatg	ctgttggtgt	780
15	atcctactgg	aaagcttctg	agggagctga	atatgatgat	cagaccagtc	aaagggagaa	840
	agaagatgat	aaagtcttcc	ctggtggaag	ccatacatat	gtctggcagg	tcctgaaaga	900
	gaatggtcca	atggcctctg	accactgtg	ccttacctac	tcatatcttt	ctcatgtgga	960
	cctggtaaaa	gacttgaatt	caggcctcat	tggagcccta	ctagtatgta	gagaagggag	1020
	tctggccaag	gaaaagacac	agaccttgca	caaatttata	ctactttttg	ctgtatttga	1080
20	tgaagggaaa	agttggcact	cagaaacaaa	gaactccttg	atgcaggata	gggatgctgc	1140
	atctgctcgg	gcctggccta	aaatgcacac	agtcaatggt	tatgtaaaca	ggtctctgcc	1200
	aggtctgatt	ggatgccaca	ggaaatcagt	ctattggcat	gtgattgga	tgggcaccac	1260
	tcctgaagt	cactcaatat	tcctcgaagg	tcacacattt	cttgtagga	accatcgcca	1320
	ggcgtccttg	gaaatctcgc	caataacttt	ccttactgct	caaacactct	tgatggacct	1380
25	tggacagttt	ctactgtttt	gtcatatctc	ttcccaccaa	catgatggca	tggaagctta	1440
	tgtcaaaagta	gacagctgtc	cagaggaacc	ccaactacga	atgaaaaata	atgaagaagc	1500
	ggaagactat	gatgatgatc	ttactgattc	tgaatggat	gtggtcaggt	ttgatgatga	1560
	caactctcct	tcctttatcc	aaattcgctc	agttgccaa	aagcatccta	aaacttgggt	1620
	acattacatt	gctgctgaag	aggaggactg	ggactatgct	cccttagtcc	tcgccccga	1680
30	tgcacagaagt	tataaaagtc	aatatttgaa	caatggccct	cagcggattg	gtaggaagta	1740
	caaaaaagtc	cgatttatgg	catacacaga	tgaaccttt	aagactcgtg	aagctattca	1800
	gcatgaatca	ggaatcttgg	gacctttact	ttatggggaa	gttgagaca	cactgttgat	1860
	tatatttaag	aatcaagcaa	gcagaccata	taacatctac	cctcacggaa	tcactgatgt	1920
	ccgtcctttg	tattcaagga	gattaccaaaa	aggtgtaaaa	catttgaagg	attttccaat	1980
35	tctgccagga	gaaatattca	aatataaatg	gacagtgact	gtagaagatg	ggccaactaa	2040
	atcagatcct	cggtgcctga	cccgtatta	ctctagtttc	gttaatatgg	agagagatct	2100
	agcttcagga	ctcattggcc	ctctcctcat	ctgtacaaa	gaatctgtag	atcaaagagg	2160
	aaaccagata	atgtcagaca	agaggaatgt	catcctgttt	tctgtatttg	atgagaaccg	2220
	aagctggtac	ctcacagaga	atatacaacg	ctttctcccc	aatccagctg	gagtgcagct	2280
40	tgaggatcca	gagttccaag	cctccaacat	catgcacagc	atcaatggct	atgtttttga	2340
	tagtttgag	ttgtcagttt	gtttgcatga	ggtggcatac	tggtagattc	taagcattgg	2400
	agcacagact	gacttccttt	ctgtcttctt	ctctggatat	accttcaaac	acaaaatggt	2460
	ctatgaagac	acactcacc	tattcccatt	ctcaggagaa	actgtcttca	tgtcgatgga	2520
	aaaccagggt	ctatggattc	tgggtgcca	caactcagac	tttcggaaca	gaggcatgac	2580
	cgccttactg	aaggtttcta	gttgtgacaa	gaactcgggt	gattattacg	aggacagtta	2640
45	tgaagatatt	tcagcatact	tgtgtagtaa	aaacaatgcc	attgaaccaa	gaagcttctc	2700
	ccagaatcca	ccagtcttga	aacgccatca	acgcgaaata	actcgtacta	ctcttcagtc	2760
	agatcaagag	gaaattgact	atgatgatac	catatcagtt	gaaatgaaga	aggaagattt	2820
	tgcattttat	gatgaggatg	aaaatcagag	ccccgcgagc	tttcaaaaaga	aaacacgaca	2880

ctatatttatt gctgcagtgg agaggctctg ggattatggg atgagtagct cccacatgt 2940
 tctaagaaac agggctcaga gtggcagtgt ccctcagttc aagaaagttg ttttccagga 3000
 atttactgat ggctccttta ctacagccctt ataccgtgga gaactaaatg aacatttggg 3060
 actcctgggg ccatatataa gagcagaagt tgaagataat atcatggtaa ctttcagaaa 3120
 5 tcaggcctct cgtccctatt ccttctattc tagccttatt tottatgagg aagatcagag 3180
 gcaaggagca gaacctagaa aaaactttgt caagcctaata gaaacaaaaa cttacttttg 3240
 gaaagtgcaa catcatatgg caccactaa agatgagttt gactgcaaag cctgggctta 3300
 tttctctgat gttgacctgg aaaaagatgt gcaactcaggc ctgattggac cccttctggt 3360
 ctgccacact aacacactga accctgctca tgggagacaa gtgacagtac aggaatttgc 3420
 10 tctgtttttc accatctttg atgagaccaa aagctggtac ttcactgaaa atatggaaag 3480
 aaactgcagg gctccctgca atatccagat ggaagatccc actttttaaag agaattatcg 3540
 cttccatgca atcaatggct acataatgga tacactacct ggcttagtaa tggctcagga 3600
 tcaaaggatt cgatggtatc tgctcagcat gggcagcaat gaaaacatcc attctattca 3660
 tttcagtggga catgtgttca ctgtacgaaa aaaagaggag tataaaatgg cactgtacaa 3720
 15 tctctatcca ggtgtttttg agacagtgga aatgttacca tccaaagctg gaatttggcg 3780
 ggtggaatgc cttattggcg agcatctaca tgctgggatg agcacacttt ttctggtgta 3840
 cagcaataag tgtcagactc ccctgggaat ggcttctgga cacattagag attttcagat 3900
 tacagcttca ggacaatatg gacagtgggc cccaaagctg gccagacttc attattccgg 3960
 atcaatcaat gcctggagca ccaaggagcc cttttcttgg atcaagggtg atctgttggc 4020
 20 accaatgatt attcacggca tcaagacca gggtgccgt cagaagtctt ccagcctcta 4080
 catctctcag tttatcatca tgtatagtct tgatgggaag aagtggcaga cttatcgagg 4140
 aaattccact ggaaccttaa tggctcttct tggcaatgtg gattcatctg ggataaaaaca 4200
 caatattttt aaccctccaa ttattgctcg atacatccgt ttgcacccaa ctcattatag 4260
 cattcgagc actcttcgca tggagttgat gggctgtgat ttaaatagtt gcagcatgcc 4320
 25 attgggaatg gagagtaaag caatatcaga tgcacagatt actgcttcat cctactttac 4380
 caatatgttt gccacctggt ctcttcaaa agctcgactt cacctccaag ggaggagtaa 4440
 tgcctggaga cctcaggtga ataatccaaa agagtggctg caagtggact tccagaagac 4500
 aatgaaagtc acaggagtaa ctactcagg agtaaaatct ctgcttacca gcatgtatgt 4560
 gaaggagttc ctcatctcca gcagtcaaga tggccatcag tggactctct tttttcagaa 4620
 30 tggcaaaagta aagggttttc agggaaatca agactccttc acacctgtgg tgaactctct 4680
 agaccaccg ttactgactc gctacctcg aattcacccc cagagttggg tgcaccagat 4740
 tgccctgagg atggaggttc tgggctgcca ggcacaggac ctctactgac tcgagcctaa 4800
 taaaggaaat ttattttcat tgcaatagtg tgttggtttt ttgtgtgctg ccgcaggaac 4860
 ccctagtgat ggagttggcc actccctctc tgcgcgctcg ctgcctcact gaggccgggc 4920
 35 gaccaaaggc cggccgacgc cggggctttg cccggcgggc ctcagtgagc gagcgagcgc 4980
 gcagctgcct gcaggacat 4999

<210> 15

<211> 14

40 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

45

<400> 15

Ser Phe Ser Gln Asn Pro Pro Val Leu Lys Arg His Gln Arg

1

5

10

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising a recombinant adeno-associated virus (AAV) virion comprising a nucleotide sequence encoding a functional Factor VIII protein, wherein said recombinant adeno-associated virus virion lacks AAV rep and cap genes, and wherein said Factor VIII protein is B domain-deleted.
2. The pharmaceutical composition of Claim 1, wherein said nucleotide sequence encoding a functional Factor VIII protein is operably linked to a tissue-specific promoter.
3. The pharmaceutical composition of Claim 1 wherein said tissue-specific promoter comprises a liver-specific promoter.
4. The pharmaceutical composition of Claim 3, wherein said promoter comprises the HNF-3 albumin promoter.
5. The pharmaceutical composition of Claim 3, wherein said promoter comprises at least a portion of the transthyretin (TTR) gene promoter.
6. The pharmaceutical composition of Claim 1 wherein said nucleotide sequence comprises a heavy chain and a light chain, and wherein said nucleotide sequence further comprises a junction that operably links said heavy chain and said light chain.
7. The pharmaceutical composition of Claim 6, wherein said junction has the amino acid sequence Ser-Phe.
8. The pharmaceutical composition of Claim 6, wherein said junction has the amino acid sequence of SEQ ID NO:15.
9. The pharmaceutical composition of Claim 1, wherein said nucleotide sequence comprises SEQ ID NO:13.

10. The pharmaceutical composition of Claim 1, wherein said nucleotide sequence comprises SEQ ID NO:14.

11. The pharmaceutical composition of Claim 1 further comprising a pharmaceutically acceptable excipient.

ABSTRACT

One form of a composition has two types of recombinant adeno-associated virus. The first type encodes a portion of Factor VIII operably linked to an expression control element; and the second type encodes a different portion of Factor VIII operably linked to an expression control element. The first and second nucleotide sequences collectively encode a functional Factor VIII protein. Another form of the composition is a recombinant adeno-associated virus containing a nucleotide sequence encoding functional Factor VIII light or heavy chain operably linked to a tissue-specific promoter.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☒ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.